PPPL-3448 UC-70

PPPL-3448

Macroscopic Description of Pressure-anisotropy-driven Collective Instability in Intense Charged Particle Beams

by Sean Strasburg and Ronald C. Davidson

June 2000



PPPL Reports Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warrantv. express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its by recommendation, or favoring endorsement, the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Availability

This report is posted on the U.S. Department of Energy's Princeton Plasma Physics Laboratory Publications and Reports web site in Calendar Year 2000. The home page for PPPL Reports and Publications is: http://www.pppl.gov/pub_report/

DOE and DOE Contractors can obtain copies of this report from:

U.S. Department of Energy Office of Scientific and Technical Information DOE Technical Information Services (DTIS) P.O. Box 62 Oak Ridge, TN 37831

Telephone: (865) 576-8401 Fax: (865) 576-5728 Email: reports@adonis.osti.gov

This report is available to the general public from:

National Technical Information Service U.S. Department of Commerce 5285 Port Royal Road Springfield, VA 22161 Telephone: 1-800-553-6847 or (703) 605-6000 Fax: (703) 321-8547 Internet: http://www.ntis.gov/ordering.htm

Macroscopic Description of Pressure-Anisotropy-Driven Collective Instability in Intense Charged Particle Beams

Sean Strasburg and Ronald C. Davidson

Plasma Physics Laboratory Princeton University Princeton, New Jersey, 08543

Abstract

The macroscopic warm-fluid model developed by Lund and Davidson [Phys. Plasmas 5, 3028 (1998)] is used in the smooth-focusing approximation to investigate detailed stability properties of an intense charged particle beam with pressure anisotropy, assuming small-amplitude electrostatic perturbations about a waterbag equilibrium. Stability properties are calculated numerically for the case of extreme pressure anisotropy with $P_{\parallel}^{0}(r) = 0$ and $P_{\perp}^{0}(r) \neq 0$, assuming axisymmetric wave perturbations $(\partial/\partial\theta = 0)$ of the form $\delta\phi(\mathbf{x},t) = \delta\hat{\phi}(r) \exp(ik_{z}z - i\omega t)$, where k_{z} is the axial wavenumber, and $Im\omega > 0$ corresponds to instability (temporal growth). For sufficiently large values of $k_{z}r_{b}$, where r_{b} is the beam radius, the analysis leads to an anisotropy-driven instability ($Im\omega > 0$) provided the normalized Debye length ($\Gamma_{D} = \lambda_{D\perp}/r_{b}$) is sufficiently large and the normalized beam intensity ($s_{b} = \hat{\omega}_{pb}^{2}/2\gamma_{b}^{2}\omega_{\beta\perp}^{2}$) is sufficiently below the space-charge limit.

I. INTRODUCTION

There is increasing interest in the equilibrium and stability properties of intense charged particle beams¹⁻³, with potential applications including heavy ion fusion, transmutation of radioactive waste, accelerator-based production of tritium, and spallation neutron sources. At the beam intensities of practical interest, it is particularly important to develop an improved theoretical understanding of the influence of space-charge effects and collective processes on detailed stability and transport properties⁴⁻²⁰. A complete description of collective processes in intense nonneutral beams generally requires a knowledge of the beam distribution function $f(\mathbf{x}, \mathbf{p}, t)$ in the six-dimensional phase space (\mathbf{x}, \mathbf{p}) . While considerable progress has been made in analytical investigations based on the Vlasov-Maxwell equations⁴⁻¹⁷, such kinetic analyses are often complex, even under idealized assumptions. It is therefore important to develop and test the robustness of alternative theoretical models, such as macroscopic models²¹⁻²⁶ based on the fluid-Maxwell equations, for investigating beam equilibrium and stability properties.

Macroscopic fluid descriptions^{21–26} have met with recent success in describing the propagation of space-charge-dominated (low-emittance) beams in periodic-focusing transport systems^{22,25}, and in describing high-frequency collective oscillations in high-intensity beams²⁴. In this Letter, we make use of the macroscopic warm-fluid model developed by Lund and Davidson²⁴ in the smooth-focusing approximation to investigate the linear stability properties of an intense charged particle beam, allowing for equilibrium pressure anisotropy $(P_{\perp}^{0} \neq P_{\parallel}^{0})$. A particular focus in the present analysis is application of the warm-fluid model to investigate the anisotropy-driven $(P_{\perp}^{0} > P_{\parallel}^{0})$ instability observed by Lund, *et al.*^{18–20} in particle-in-cell simulations and studied analytically using the Vlasov-Maxwell equations. Such anisotropies are well known to develop naturally in accelerators. For example, for a beam of charged particles of mass *m* and charge *q* that is accelerated through a voltage *V*, a simple estimate shows that the final and initial longitudinal temperatures (in energy units) are related⁴ in the nonrelativistic case by $T_{\parallel f} = T_{\parallel f}^{2}/2qV$. In the relativistic case, this

relation is modified to become $T_{\parallel f} = T_{\parallel i}^2 \gamma_i^3 / \beta_f^2 \gamma_j^3 mc^2$, where γ is the relativistic mass factor and β is the relativistic velocity. As an example, for an electron beam with initial energy 10 keV and temperature $T_{\parallel i} = 0.5$ eV accelerated to 1 MeV, the final longitudinal temperature is $T_{\parallel f} = 2.1 \times 10^{-8}$ eV, a decrease by seven orders-of-magnitude. In addition, the beam's effective transverse temperature T_{\perp} and emittance are subject to increase due to nonlinearities in applied and self-field forces, nonstationary beam profiles, and mismatches, which may produce negligible changes in the parallel temperature. This simultaneous cooling in the parallel direction and heating in the transverse direction can provide the free energy to drive collective instabilities and cause a further deterioration in beam quality through the instability mechanism described in this paper. As a general remark, application of a warmfluid model to describe the equilibrium and stability properties of intense charged particle beams appears to be a remarkably robust and simple approach, both for the case of stable high-frequency collective oscillations²⁴, as well as the unstable case considered here, where the instability is driven by gross macroscopic properties of the beam equilibrium (pressure anisotropy).

To briefly summarize the assumptions and macroscopic warm-fluid model, the present analysis considers an intense nonneutral beam consisting of charged particles with charge q and rest mass m propagating in the z-direction with average axial velocity $V_b = const.$, and characteristic directed kinetic energy $(\gamma_b - 1)mc^2$ in the laboratory frame. Here, $\gamma_b =$ $(1 - V_b^2/c^2)^{-1/2}$ is the relativistic mass factor, c is the speed of light *in vacuo*, and a perfectly conducting cylindrical wall is located at radius $r = r_w$, where $r = (x^2 + y^2)^{1/2}$ is the radial distance from the beam axis. The characteristic beam radius is denoted by r_b , and it is assumed that the particle motion in the beam frame is nonrelativistic. Transverse confinement of the beam particles is provided by applied magnetic or electric focusing fields, and in the *smooth-focusing* approximation we model the applied transverse focusing force on a beam particle by $\mathbf{F}_{foc} = -\gamma_b m \omega_{\beta\perp}^2 (x \hat{\mathbf{e}}_x + y \hat{\mathbf{e}}_y)$, where $\omega_{\beta\perp} = const$. is the effective betatron frequency for the transverse oscillations, and (x, y) is the transverse displacement from the beam axis. The present analysis is carried out in the electrostatic approximation, where the selfelectric field $\mathbf{E}^{s}(\mathbf{x}, t)$ produced by the beam space charge is $\mathbf{E}^{s} = -\nabla\phi$, and the electrostatic potential $\phi(\mathbf{x}, t)$ is determined self-consistently from Poisson's equation, $\nabla^{2}\phi = -4\pi q n(\mathbf{x}, t)$, where $n(\mathbf{x}, t)$ is the number density. In addition, the axial beam current, $qn(\mathbf{x}, t)V_{z}(\mathbf{x}, t)$, where $V_{z}(\mathbf{x}, t)$ is the average axial velocity, produces a transverse self-magnetic field, $\mathbf{B}^{s}(\mathbf{x}, t)$, where $\mathbf{B}^{s} = \nabla A_{z} \times \hat{\mathbf{e}}_{z}$. Here, $A_{z}(\mathbf{x}, t)$ is determined self-consistently in the magnetostatic approximation from Maxwell's equation, $\nabla^{2}A_{z} = -(4\pi/c)qnV_{z}$. In circumstances where the average axial velocity is approximately uniform over the beam cross section with $V_{z} \simeq V_{b} =$ const., which we assume to be the case, the self-field potentials, $\phi(\mathbf{x}, t)$ and $A_{z}(\mathbf{x}, t)$, are related by the familiar expression, $A_{z} = (V_{b}/c)\phi$.

To describe the dynamics of the intense charged particle beam interacting with the applied focusing field and the self-generated electric and magnetic fields, we make use of the macroscopic warm-fluid model developed by Lund and Davidson²⁴. To briefly summarize, by taking appropriate momentum moments of the nonlinear Vlasov equation for the beam distribution function $f(\mathbf{x}, \mathbf{p}, t)$, we obtain an interconnected chain of macroscopic fluid equations²¹ advancing the particle density $n(\mathbf{x}, t)$, the average flow velocity $\mathbf{V}(\mathbf{x}, t) = V_z(\mathbf{x}, t)\hat{\mathbf{e}}_z + \mathbf{V}_{\perp}(\mathbf{x}, t)$, the pressure tensor $\mathbf{P}(\mathbf{x}, t)$, the heat flow tensor $\mathbf{Q}(\mathbf{x}, t)$, etc. Following Lund and Davidson²⁴, we adopt a model in which the heat-flow contribution, proportional to $(\partial/\partial \mathbf{x}) \cdot \mathbf{Q}(\mathbf{x}, t)$, is neglected in the dynamical equation advancing the pressure tensor $\mathbf{P}(\mathbf{x}, t)$. In addition, we adopt a model²¹ in which $\mathbf{P}(\mathbf{x}, t)$ is assumed to be isotropic in the plane perpendicular to the beam propagation direction (the z-direction), i.e., $\mathbf{P}(\mathbf{x}, t) = P_{\perp}(\mathbf{x}, t)(\hat{\mathbf{e}}_x \hat{\mathbf{e}}_x + \hat{\mathbf{e}}_y \hat{\mathbf{e}}_y) + P_{\parallel}(\mathbf{x}, t)\hat{\mathbf{e}}_z \hat{\mathbf{e}}_z$, where $P_{\perp}(\mathbf{x}, t)$ and $P_{\parallel}(\mathbf{x}, t)$ are scalar pressures. In this case, making use of the assumptions enumerated earlier, the warm-fluid equations²⁴ appropriately generalized to the case of relativistic axial motion are given by the continuity equation for $n(\mathbf{x}, t)$,

$$\left(\frac{\partial}{\partial t} + V_z \frac{\partial}{\partial z} + \mathbf{V}_\perp \cdot \frac{\partial}{\partial \mathbf{x}_\perp}\right) n + n \left(\frac{\partial V_z}{\partial z} + \frac{\partial}{\partial \mathbf{x}_\perp} \cdot \mathbf{V}_\perp\right) = 0 , \qquad (1)$$

the perpendicular force balance equation for $\mathbf{V}_{\perp}(\mathbf{x}, t)$,

$$\gamma_b mn \left(\frac{\partial}{\partial t} + V_z \frac{\partial}{\partial z} + \mathbf{V}_\perp \cdot \frac{\partial}{\partial \mathbf{x}_\perp} \right) \mathbf{V}_\perp + \frac{\partial}{\partial \mathbf{x}_\perp} P_\perp = -nq \frac{1}{\gamma_b^2} \nabla_\perp \phi - \gamma_b mn \omega_{\beta\perp}^2 \mathbf{x}_\perp , \qquad (2)$$

the parallel force balance equation for $V_z(\mathbf{x}, t)$,

$$\gamma_b mn \left(\frac{\partial}{\partial t} + V_z \frac{\partial}{\partial z} + \mathbf{V}_\perp \cdot \frac{\partial}{\partial \mathbf{x}_\perp} \right) V_z + \frac{\partial}{\partial z} P_{||} = -nq \frac{\partial \phi}{\partial z} , \qquad (3)$$

the equation of state for the perpendicular pressure $P_{\perp}(\mathbf{x}, t)$,

$$\left(\frac{\partial}{\partial t} + V_z \frac{\partial}{\partial z} + \mathbf{V}_\perp \cdot \frac{\partial}{\partial \mathbf{x}_\perp}\right) \left(\frac{P_\perp}{n^2}\right) - \frac{P_\perp}{n^2} \frac{\partial V_z}{\partial z} = 0 , \qquad (4)$$

the equation of state for the parallel pressure $P_{\parallel}(\mathbf{x}, t)$,

$$\left(\frac{\partial}{\partial t} + V_z \frac{\partial}{\partial z} + \mathbf{V}_\perp \cdot \frac{\partial}{\partial \mathbf{x}_\perp}\right) \left(\frac{P_{\parallel}}{n}\right) + \frac{2P_{\parallel}}{n} \frac{\partial V_z}{\partial z} = 0 , \qquad (5)$$

and Poisson's equation for the electrostatic potential $\phi(\mathbf{x}, t)$,

$$\nabla_{\perp}^{2}\phi + \frac{\partial^{2}}{\partial z^{2}}\phi = -4\pi qn \ . \tag{6}$$

Equations (1)–(6) provide a closed macroscopic description of the nonlinear evolution of the beam interacting with the applied focusing field and the self-generated electric and magnetic fields, $\mathbf{E}^s = -\nabla\phi$ and $\mathbf{B}^s = (V_b/c)\nabla\phi \times \hat{\mathbf{e}}_z$. In obtaining Eqs. (1)–(6), it has been assumed that the fluid motions in the beam frame are nonrelativistic, i.e., \mathbf{V}_{\perp}^2/c^2 , $(V_z - V_b)^2/c^2$, $P_{\perp}/\gamma_b nmc^2$, $P_{\parallel}/\gamma_b nmc^2 \ll 1$.

Equations (1)–(6) can be used to investigate detailed macroscopic equilibrium and stability properties for perturbations about a wide range of beam equilibria²⁴ ranging from a warmfluid thermal equilibrium with diffuse radial density profile, to a warm-fluid Kapchinskij-Vladimirskij (KV) equilibrium with step-function density profile, to a warm-fluid waterbag equilibrium. For example, assuming $\partial/\partial z = 0$, Lund and Davidson have investigated²⁴ stable electrostatic oscillations for perturbations about a warm-fluid KV equilibrium. In the present analysis, allowing for a pressure anisotropy with $P_{\perp} \neq P_{\parallel}$ and perturbations with $\partial/\partial z \neq 0$, we examine detailed stability properties for perturbations about a warm-fluid waterbag equilibrium^{11,13,24}. Under steady-state (equilibrium) conditions with $\partial/\partial t = 0$, we assume a matched, axisymmetric beam in which all equilibrium profiles (denoted with a superscript zero), satisfy $\partial/\partial \theta = 0 = \partial/\partial z = 0$, and depend only on the radial distance $r = (x^2 + y^2)^{1/2}$ from the beam axis. In equilibrium, it is further assumed that there is no perpendicular motion of the beam and that the axial flow velocity is uniform over the beam cross section, i.e., $\mathbf{V}^0_{\perp} = 0$, and $V_z^0 = V_b = const$. For a warm-fluid waterbag equilibrium^{11,13,24}, we assume that the perpendicular and parallel pressures are of the form

$$P^{0}_{\perp}(r) = (\hat{T}_{\perp}/\hat{n})[n^{0}(r)]^{2} , \qquad P^{0}_{\parallel}(r) = \hat{T}_{\parallel}n^{0}(r) , \qquad (7)$$

where $n^0(r)$ is the equilibrium density profile. In Eq. (7), $\hat{n} = n^0(r = 0) = const$. is the on-axis density, and the constants \hat{T}_{\perp} and \hat{T}_{\parallel} are the perpendicular temperature and parallel temperature, respectively, at r = 0, expressed in energy units. From Eq. (7), we note that the effective temperature profiles, $T^0_{\perp}(r) = P^0_{\perp}(r)/n^0(r)$ and $T^0_{\parallel}(r) = P^0_{\parallel}(r)/n^0(r)$, are given by $T^0_{\perp}(r) = \hat{T}_{\perp}n^0(r)/\hat{n}$ and $T^0_{\parallel}(r) = \hat{T}_{\parallel} = const$. That is, $T^0_{\perp}(r)$ has the same radial shape as the density profile $n^0(r)$, whereas $T^0_{\parallel}(r)$ is uniform (isothermal) over the beam cross section. We introduce the effective perpendicular Debye length $\lambda_{D\perp}$ and self-field intensity parameter s_b defined by

$$\lambda_{D\perp}^{2} = \frac{2\hat{T}_{\perp}\gamma_{b}^{2}}{4\pi q^{2}\hat{n}} , \qquad s_{b} = \frac{\hat{\omega}_{pb}^{2}}{2\gamma_{b}^{2}\omega_{\beta\perp}^{2}} , \qquad (8)$$

where $\hat{\omega}_{pb}^2 = 4\pi \hat{n}q^2/\gamma_b m$ is the on-axis plasma frequency-squared. Without presenting algebraic details, the exact solution for the equilibrium density profile is given by²⁶

$$n^{0}(r) = \begin{cases} \hat{n} \frac{I_{0}(r_{b}/\lambda_{D\perp}) - I_{0}(r/\lambda_{D\perp})}{I_{0}(r_{b}/\lambda_{D\perp}) - 1} , & 0 \leq r < r_{b} , \\ 0 , & r_{b} < r \leq r_{w} , \end{cases}$$
(9)

where r_b is the outer radius of the beam, and r_w is the conducting wall radius. In terms of $\lambda_{D\perp}$ and $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2 \omega_{\beta\perp}^2$, the beam radius r_b in Eq. (9) is determined self-consistently from

$$I_0(r_b/\lambda_{D\perp}) = \frac{1}{1 - \hat{\omega}_{pb}^2/2\gamma_b^2 \omega_{\beta\perp}^2} .$$
 (10)

Here, $I_0(x)$ is the modified Bessel function of the first kind of order zero.

For the equilibrium density profile $n^0(r)$ specified by Eq. (9), we note that the density profile decreases monotonically from the on-axis value $n^0(r=0) = \hat{n}$ at r=0, to $n^0(r=r_b) = 0$ at the beam edge $(r=r_b)$. At low beam intensities with $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2 \omega_{\beta\perp}^2 \ll 1$, it follows from Eqs. (9) and (10) that $r_b < \lambda_{D\perp}$, corresponding to an emittance-dominated beam with near-parabolic density profile, $n^0(r) = \hat{n}(1 - r^2/r_b^2)$, over the beam cross section. On the other hand, for $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2 \omega_{\beta\perp}^2 \to 1 - \epsilon$, with $\epsilon \to 0_+$, it follows from Eqs. (9) and (10) that $r_b \gg \lambda_{D\perp}$ and that $n^0(r)$ approaches the step-function density profile characteristic of space-charge-dominated beams with very low transverse emittance. These properties are illustrated in Fig. 1, where the normalized density profile $n^0(r)/\hat{n}$ calculated from Eq. (9) is plotted versus r/r_b for several values of the dimensionless intensity parameter $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2 \omega_{\beta\perp}^2$ over the interval $0 < s_b < 1$.

The macroscopic fluid-Poisson equations (1)–(6) can be linearized for small-amplitude perturbations about the warm-fluid waterbag equilibrium described by Eqs. (7), (9), $\mathbf{V}^{0}_{\perp}(\mathbf{x}) = 0$, and $V_{z}^{0} = V_{b} = const$. The resulting linearized equations then can be combined into a single eigenvalue equation²⁶ for the perturbed electrostatic potential $\delta\phi(\mathbf{x}, t)$, allowing for arbitrary anisotropy in the perpendicular and parallel pressures, $P^{0}_{\perp}(r)$ and $P^{0}_{\parallel}(r)$. The details of this derivation will be presented elsewhere²⁶.

Using a normal-mode approach, we express all perturbed quantities $\delta \phi(\mathbf{x}, t)$, $\delta P_{\perp}(\mathbf{x}, t)$, $\delta P_{\parallel}(\mathbf{x}, t)$, and $\delta n(\mathbf{x}, t)$ as

$$\delta\psi(\mathbf{x},t) = \delta\hat{\psi}(\mathbf{x}_{\perp})\exp(ik_z z - i\omega t) .$$
⁽¹¹⁾

Here, $\delta \hat{\psi}(\mathbf{x}_{\perp})$ is the perturbation amplitude, k_z is the axial wavenumber of the perturbation, and ω is the complex oscillation frequency, with $Im\omega > 0$ corresponding to instability (temporal growth). We further introduce the definitions

$$\Omega = \omega - k_z V_b , \quad \omega_{pb}^2(r) = \frac{4\pi n^0(r)q^2}{\gamma_b m} , \quad v_{Tz}^2 = \frac{2\tilde{T}_{||}}{\gamma_b m} , \qquad (12)$$

where Ω is the Doppler-shifted oscillation frequency, $\omega_{pb}^2(r)$ is the local relativistic plasma

frequency-squared, and $v_{Tz} = const.$ is the effective axial thermal speed of the beam ions. After some algebraic manipulation and rearrangement of terms, we obtain²⁶

$$\frac{\partial}{\partial \mathbf{x}_{\perp}} \cdot \left\{ \left[\Omega^{2} - \frac{\omega_{pb}^{2}(r)}{\gamma_{b}^{2}} (1 + k_{z}^{2} \lambda_{D\perp}^{2}) - \frac{\frac{1}{2} k_{z}^{2} v_{TZ}^{2} \Omega^{2}}{\Omega^{2} - k_{z}^{2} v_{TZ}^{2}} \right] \frac{\partial}{\partial \mathbf{x}_{\perp}} \delta \hat{\phi}
+ \lambda_{D\perp}^{2} \frac{\omega_{pb}^{2}(r)}{\gamma_{b}^{2}} \frac{\partial}{\partial \mathbf{x}_{\perp}} \nabla_{\perp}^{2} \delta \hat{\phi}
+ \frac{1}{2} k_{z}^{2} \lambda_{D\perp}^{2} \frac{\partial}{\partial \mathbf{x}_{\perp}} \left[\frac{\omega_{pb}^{2}(r) / \gamma_{b}^{2}}{\Omega^{2} - k_{z}^{2} v_{TZ}^{2}} \left([\omega_{pb}^{2}(r) + \frac{1}{2} k_{z}^{2} v_{TZ}^{2}] \delta \hat{\phi} - \frac{1}{2} v_{TZ}^{2} \nabla_{\perp}^{2} \delta \hat{\phi} \right) \right] \right\}
- k_{z}^{2} \left\{ \Omega^{2} - \frac{\omega_{pb}^{2}(r) \Omega^{2}}{\Omega^{2} - k_{z}^{2} v_{TZ}^{2}} - \frac{\frac{1}{2} k_{z}^{2} v_{TZ}^{2} \Omega^{2}}{\Omega^{2} - k_{z}^{2} v_{TZ}^{2}} \right\} \delta \hat{\phi} = 0.$$
(13)

Equation (13) is the final eigenvalue equation for the potential eigenfunction $\delta \hat{\phi}(\mathbf{x}_{\perp})$ and the eigenfrequency $\Omega = \omega - k_z V_b$. Equation (13) has been derived from Eqs. (1)–(6) for smallamplitude perturbations about the warm-fluid waterbag equilibrium described by Eqs. (7) and (9), and can be used to investigate detailed stability properties over a wide range of values of beam intensity and temperature anisotropy. Introducing cylindrical polar coordinates (r, θ) , where $x = r \cos \theta$ and $y = r \sin \theta$, we represent $\delta \hat{\phi}(\mathbf{x}_{\perp}) = \delta \hat{\phi}(r, \theta)$. The eigenvalue equation (13) is to be solved for $\delta \hat{\phi}(r, \theta)$ and Ω subject to the requirements that $\delta \hat{\phi}(r, \theta)$ be regular at the origin (r = 0), and that

$$\delta\hat{\phi}(r = r_w , \theta) = 0 . \tag{14}$$

The boundary condition in Eq. (14) assures that the perturbed tangential electric field components vanish at the perfectly conducting wall.

Anisotropy-driven instabilities are expected to be the strongest (largest growth rate) in the case of strong temperature anisotropy $\hat{T}_{\perp} \gg \hat{T}_{\parallel}$. In circumstances where the beam ions are cold in the propagation direction, the eigenvalue equation (13) simplifies considerably. Therefore, in the remainder of this article we set $\hat{T}_{\parallel} = 0$ ($v_{Tz}^2 = 0$) in Eq. (13), and further assume axisymmetric perturbations with $(\partial/\partial\theta)\delta\hat{\phi} = 0$. The eigenvalue equation (13) then reduces to²⁶

$$\frac{1}{r} \frac{\partial}{\partial r} r \left\{ \left[\Omega^2 - \frac{\omega_{pb}^2(r)}{\gamma_b^2} (1 + k_z^2 \lambda_{D\perp}^2) \right] \frac{\partial}{\partial r} \delta \hat{\phi}(r) + \lambda_{D\perp}^2 \frac{\omega_{pb}^2(r)}{\gamma_b^2} \left(\frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} - \frac{1}{r^2} \right) \frac{\partial}{\partial r} \delta \hat{\phi}(r) + \frac{1}{2} k_z^2 \lambda_{D\perp}^2 \frac{\partial}{\partial r} \left(\frac{\omega_{pb}^4(r)/\gamma_b^2}{\Omega^2} \delta \hat{\phi}(r) \right) \right\} - k_z^2 [\Omega^2 - \omega_{pb}^2(r)] \delta \hat{\phi}(r) = 0 .$$
(15)

For $k_z = 0$ and $\lambda_{D\perp} \neq 0$, it is found²⁶ that Eq. (15) gives purely stable oscillations with $Im\hat{\Omega} = 0$. On the other hand, as $k_z r_b$ is increased to sufficiently large values, the temperature anisotropy $(\hat{T}_{\perp} \propto \lambda_{D\perp}^2 \neq 0$, and $\hat{T}_{\parallel} = 0)$ provides the free energy to drive an instability²⁶ at moderate values of beam intensity.

For the equilibrium density profile $n^0(r)$ specified by Eq. (9), the eigenvalue equation (15) must be solved numerically for the (generally) complex eigenfunction $\delta \hat{\phi} = \delta \hat{\phi}_r + i\delta \hat{\phi}_i$ and complex eigenfrequency $\Omega = \Omega_r + i\Omega_i = \omega_r + i\omega_i - k_z V_b$, subject to the boundary condition at $r = r_w$ in Eq. (14). A detailed investigation of Eq. (15) for a wide range of system parameters will be presented elsewhere²⁶, and we summarize here several of the key results. First, the dimensionless parameters characterizing Eq. (15) are

$$k_z r_b , \quad \Gamma_D \equiv \frac{\lambda_{D\perp}}{r_b} , \quad s_b \equiv \frac{\omega_{pb}^2}{2\gamma_b^2 \omega_{\beta\perp}^2} , \quad \frac{r_b}{r_w} ,$$
 (16)

where s_b and Γ_D are related by Eq. (10). Second, it is convenient to introduce the characteristic measures of the depressed and undepressed single-particle transverse oscillation frequencies, ν and ν_0 , defined by

$$\nu^2 \equiv \omega_{\beta\perp}^2 - \hat{\omega}_{pb}^2 / 2\gamma_b^2 \ , \quad \nu_0^2 \equiv \omega_{\beta\perp}^2 \ , \tag{17}$$

and the effective tune depression, ν/ν_0 , where

$$\frac{\nu^2}{\nu_0^2} = 1 - \frac{\hat{\omega}_{pb}^2}{2\gamma_b^2 \omega_{\beta\perp}^2} = 1 - s_b \ . \tag{18}$$

Finally, in Eq. (15) there is a clear demarcation between the vacuum region where $\omega_{pb}^2(r) = 0$ $(r_b < r \leq r_w)$, and the beam interior $(0 \leq r < r_b)$. For $s_b \neq 1$ and $\Gamma_D \neq 0$, the equilibrium density profile $n^0(r)$ in Eq. (9) approaches zero continuously at $r = r_b$ (see Fig. 1), and it is readily shown²⁶ from Eq. (15) that the appropriate boundary conditions at the beam surface $(r = r_b)$ correspond to continuity of $\delta \hat{\phi}(r)$,

$$Re[\delta\hat{\phi}^{I}(r=r_{b})] = Re[\delta\hat{\phi}^{II}(r=r_{b})] ,$$

$$Im[\delta\hat{\phi}^{I}(r=r_{b})] = Im[\delta\hat{\phi}^{II}(r=r_{b})] ,$$
(19)

and continuity of $(\partial/\partial r)\delta\hat{\phi}(r)$,

$$Re\left[\frac{\partial}{\partial r}\delta\hat{\phi}^{I}(r)\right]_{r=r_{b}} = Re\left[\frac{\partial}{\partial r}\delta\hat{\phi}^{II}(r)\right]_{r=r_{b}},$$

$$Im\left[\frac{\partial}{\partial r}\delta\hat{\phi}^{I}(r)\right]_{r=r_{b}} = Im\left[\frac{\partial}{\partial r}\delta\hat{\phi}^{II}(r)\right]_{r=r_{b}},$$
(20)

where Region I corresponds to the beam interior $(0 \le r < r_b)$, and Region II corresponds to the vacuum region $(r_b < r \le r_w)$.

For $k_z = 0$, it is readily shown from Eq. (15) that the radial electric field perturbation, $\delta \hat{E}_r = -(\partial/\partial r)\delta \hat{\phi}$, is equal to zero in the vacuum region, i.e., $\delta \hat{E}_r = 0$ for $r_b < r \leq r_w$. For $k_z = 0$, Eq. (15) has been solved²⁶ over the interval $0 \leq r < r_b$ for $\delta \hat{E}_r(r)$ and Ω subject to the requirement that $[\delta \hat{E}_r]_{r=r_b} = 0$ using two approaches: (a) direct numerical integration of Eq. (15) using a shooting method to determine both the eigenfunctions and eigenvalues, and (b) a matrix-dispersion-equation technique that expands Eq. (15) in a complete set of basis functions that satisfy a priori the boundary condition $[\delta \hat{E}_r]_{r=r_b} = 0$. The results using both techniques are in excellent agreement²⁶. For $k_z = 0$, the solutions to Eq. (15) correspond to a discrete set of stable oscillations $\{\omega_n\}$ with radial mode numbers n = 1, 2, 3, ..., and $Im\omega_n = 0$. Typical numerical results are shown in Fig. 2 where $Re[\omega_n/\nu_0]$ is plotted versus ν/ν_0 in Fig. 2(a) for effective tune depressions ranging from $\nu/\nu_0 = 0$ ($s_b = 1$) to $\nu/\nu_0 = 1$ ($s_b = 0$). For each value of n, note that the $\nu/\nu_0 = 0$ ($\hat{\omega}_{pb}^2/2\gamma_b^2 \hat{\omega}_{p\perp}^2 = 1$) intercept in Fig. 2(a) corresponds to the single frequency $Re\omega_n = \sqrt{2}\nu_0 = \sqrt{2}\omega_{\beta\perp} = \hat{\omega}_{pb}/\gamma_b$ in the limit of high beam intensity. On the other hand, as $\nu/\nu_0 = (1 - s_b)^{1/2}$ is increased (decreasing beam intensity $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2 \hat{\omega}_{p\perp}^2$), it is evident from Fig. 2(a) that there is a discrete spectrum of stable oscillations with frequencies $\{\omega_n\}$ that increase as ν/ν_0 and n are increased. The general features of the solutions for $\{\omega_n\}$ presented in Fig. 2(a) for $k_z = 0$ perturbations about a warm-fluid waterbag equilibrium are qualitatively similar to those for a warm-fluid Kapchinskij-Vladimirskij (KV) equilibrium²⁴, although the precise values of $\{\omega_n\}$ differ as ν/ν_0 is increased. For completeness, shown in Fig. 2(b) are plots of the eigenfunction $\delta \hat{E}(r)$ versus r/r_b for radial mode numbers n = 1, 2, 3 obtained numerically from Eq. (15) for $s_b = 0.36$ and $\nu/\nu_0 = 0.8$. Note from Fig. 2(b) that the number of radial oscillations of $\delta \hat{E}(r)$ increase as the mode number n is increased. Moreover, for specified mode number n, the number of zeros of $\delta \hat{E}(r)$ in the interval $0 \le r \le r_b$ is equal to n + 1.

For $k_z \neq 0$ and $\Gamma_D = \lambda_{D\perp}/r_b \neq 0$, the eigenvalue equation (15), valid for $\hat{T}_{\parallel} = 0$, generally supports unstable solutions $(Im\omega > 0)$ provided $k_z r_b$ is sufficiently large and the normalized beam intensity s_b is sufficiently below the space-charge limit²⁶. The eigenvalue equation (15) is a linear fourth-order ordinary differential equation for $\delta \hat{\phi}(r)$. At r = 0, several of the coefficients are singular. At $r = r_b$, the beam edge, the coefficient multiplying the highest-derivative term vanishes, causing a boundary layer. Since standard numerical integration techniques are not applicable, we instead expand the solution near r = 0 in a Froebenius series. Using this analytical expansion near the origin where it is sufficiently accurate, we begin by numerically integrating from very near r = 0 out to the beam edge at $r = r_b$, and then back to the origin.

The linearity of Eq. (15) implies that the solution for $\delta \hat{\phi}(r)$ is arbitrary up to a constant, multiplicative, complex factor. This freedom can be used in the unstable case to make the matching of real and imaginary parts at the beam edge simpler. For present purposes, we choose the initial amplitude and phase at r = 0 such that $Re[\delta \hat{\phi}(r = 0)] = 1$ and $Im[\delta \hat{\phi}(r = 0)] = 0$. In the unstable case, integrating away from r = 0, the eigenfunction generally develops an imaginary component with $Im[\delta \hat{\phi}(r)] \neq 0$. Of course at the beam edge $(r = r_b)$, there are four conditions to satisfy, corresponding to Eqs. (19) and (20). We can automatically satisfy the matching conditions in Eq. (19) by appropriate choice of complex phase factor for the solution for $\delta \hat{\phi}(r)$ in the vacuum region $(r_b < r \leq r_w)$. In general, however, neither of the matching conditions on $\partial \delta \hat{\phi} / \partial r$ at $r = r_b$ in Eq. (20) will be satisfied unless the complex eigenfrequency $\Omega = \Omega_r + i\Omega_i$ occurring in Eq. (15) is correctly chosen, which corresponds to the desired dispersion relation. Therefore, in the present shooting method, Eq. (15) is repeatedly integrated, and the value of $\Omega_r + i\Omega_i$ adjusted until the matching conditions in Eq. (19) are satisfied, thereby determining the eigenfrequency.

Using this method, Eq. (15) has been solved numerically subject to the boundary conditions in Eqs. (14), (19) and (20), and the complex eigenfrequency $\Omega = \Omega_r + i\Omega_i$ and eigenfunction $\delta \hat{\phi}(r)$ have been determined self-consistently over a wide range of system parameters corresponding to normalized beam intensity, $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2 \omega_{\beta\perp}^2$, tune depression, $\nu/\nu_0 = (1 - s_b)^{1/2}$, transverse Debye length, $\Gamma_D = \lambda_{D\perp}/r_b$, and axial wavenumber $k_z r_b$. Here, keep in mind that s_b and Γ_D are related by Eq. (10), so that very high beam intensity ($s_b \rightarrow 1$) corresponds to $\Gamma_D \ll 1$, and low beam intensity ($s_b \ll 1$) corresponds to $\Gamma_D \gg 1$ (see Fig. 1). Because $\hat{T}_{\parallel} = 0$ is assumed in the present analysis, the term proportional to $k_z^2 \lambda_{D\perp}^2 \neq 0$ in Eq. (15) provides the free energy to drive instability associated with temperature anisotropy ($\hat{T}_{\perp} > \hat{T}_{\parallel}$).

For $k_z \neq 0$, typical numerical results obtained from Eq. (15), are illustrated in Figs. 3–5 for a mildly relativistic beam with $\gamma_b = 1.02$ and $r_w/r_b = 2$. As a general remark, beams which are cold in the transverse direction oscillate stably at all values of axial wavelength. Beams with intermediate transverse temperatures are unstable for *all* wavenumbers $k_z r_b$ larger than a critical value. Finally, beams which are sufficiently hot in the transverse direction have a *finite* instability bandwidth in $k_z r_b$: short wavelengths and long wavelengths are stable, while wavelengths in a range about $k_z r_b \Gamma_D \sim 1$ are unstable. That is, the growth rate of sufficiently temperature-dominated beams turns over and approaches zero at large values of $k_z r_b$. For the choice of waterbag equilibrium considered here with $\hat{T}_{\parallel} = 0$, the onset of instability occurs for $\Gamma_D > \Gamma_D^* = 0.364$, $s_b < s_b^* = 0.750$, and $\nu/\nu_0 > \nu^*/\nu_0 = 0.500$, which are equivalent conditions. For increasing values of Γ_D relative to Γ_D^* , the instability bandwidth first increases, encompassing both higher and lower axial wavenumbers. For even warmer beams, however, high values of $k_z r_b$ become stabilized, while the region of instability continues to shift to smaller axial wavenumbers.

Figure 3 shows typical numerical results for the choice of system parameters $\Gamma_D = 0.360$ $(s_b = 0.755)$, corresponding to *stable* oscillations with $Im\Omega = Im\omega = 0$. Plotted in Fig. 3(a) is $Re(\omega - k_z V_b)/\nu_0$ versus $k_z r_b$ for the n = 1 eigenmode, whereas Fig. 3(b) shows the corresponding eigenfunction $\delta \hat{\phi}(r)$ (assumed real) plotted versus r/r_b for several values of $k_z r_b$. The high-frequency (upper) branch in Fig. 3(b) corresponds to the familiar plasma oscillation branch considered in Fig. 2, extended to non-zero values of $k_z r_b$. The lower (slow-wave) branch in Fig. 3(a) starts at zero frequency for $k_z = 0$, and $Re(\omega - k_z V_b)$ increases linearly with $k_z r_b$ for $k_z r_b < 1$, and then asymptotes at $Re(\omega - k_z V_b) \simeq 1.03\nu_0$ for $k_z r_b \gg 1$. On the other hand, the upper (high-frequency) branch in Fig. 3(a) asymptotes at $Re(\omega - k_z V_b) \simeq 1.3\nu_0$ for $k_z r_b \gg 1$. From Fig. 3(b), as expected, as $k_z r_b$ is increased, the eigenfunction $\delta \hat{\phi}(r)$ extends radially well into the vacuum region, with $\delta \hat{\phi}(r = r_w) = 0$.

For $\Gamma_D > \Gamma_D^*$, the two branches in Fig. 3 coalesce as $k_z r_b$ is increased beyond some critical value $k_z^*(\Gamma_D)r_b$. Typical numerical results in this case are illustrated in Fig. 4 for the choice of system parameters $\Gamma_D = 0.509$ ($s_b = 0.55$). From Fig. 4, for $0 \le k_z r_b <$ $k_z^* r_b = 0.968$, the eigenvalue equation supports two real oscillatory solutions with $Im\omega = 0$. For $k_b r_b > k_z^* r_b = 0.968$, however, the two modes coalesce and have the same value of $Re(\omega - k_z V_b)$, and complex conjugate values of $Im\omega$ (one mode is damped, and the other is growing). The normalized growth rate $Im\omega/\nu_0$ of the unstable branch is plotted versus $k_z r_b$ in Fig. 4(a), and increases from $Im\omega = 0$ at $k_z r_b = k_z^* r_b = 0.968$, to $Im\omega \simeq 0.4\nu_0$ for $k_z r_b \gg 1$. Consistent with Fig. 4(a), the corresponding eigenfunction plots of $Re[\delta\hat{\phi}(r)]$ and $Im[\delta\hat{\phi}(r)]$ versus r/r_b are presented in Fig. 4(b) for $k_b r_b = 4$, corresponding to instability. For moderately low values of $k_z r_b$, the eigenfunction for the unstable mode has the distinctive n = 1 mode structure illustrated in Fig. 4(b) for $k_z r_b = 4$. As $k_z r_b$ is increased, however, the real part of the eigenfunction, $Re[\delta\hat{\phi}(r)]$, changes continuously from an n = 1 to an n = 2 mode structure²⁶. For very large $k_z r_b > 10$, the boundary layer at $r = r_b$ becomes very sharp, with large changes in $Re[\delta\hat{\phi}(r)]$ over a very short radial scale²⁶. The radial mode number, however, does not appear to change from n = 2. Of course, perturbations with such large values of $k_z r_b$ are of limited practical interest because the modes would be stabilized $(Im\omega = 0)$ at short axial wavelengths by finite $\hat{T}_{\parallel} \neq 0$ effects in an analysis of the more complete eigenvalue equation (13).

For completeness, shown in Fig. 5 are plots of the normalized growth rate $Im\omega/\nu_0$ versus $k_z r_b$ obtained numerically from Eqs. (14), (15), (19), and (20) for several values of $\Gamma_D > \Gamma_D^*$ and $s_b < s_b^*$. Note from Fig. 5 that critical value of $k_z r_b$ for onset of instability increases as Γ_D is increased (s_b is decreased), and that the maximum normalized growth rate $(Im\omega)_{max}/\nu_0$ increases as Γ_D is increased (s_b is decreased). For sufficiently large value of Γ_D (large enough transverse emittance), we also note from Fig. 5 that the instability has a finite bandwidth in $k_z r_b$, whereas for smaller values of Γ_D , the maximum growth rate occurs for $k_z r_b \gg 1$. For $\hat{T}_{\parallel} \neq 0$ (but $\hat{T}_{\parallel} < \hat{T}_{\perp}$), it is expected that the more complete eigenvalue equation (13) will always give a finite instability bandwidth in $k_z r_b$.

To summarize, for $k_z = 0$, the analysis of the eigenvalue equation (15) leads to a discrete spectrum $\{\omega_n\}$ of stable oscillations with $Im\omega_n = 0$, where n is the radial mode number. On the other hand, for sufficiently large values of $k_z r_b$, where r_b is the beam radius, the analysis of the eigenvalue equation leads to an anisotropy-driven instability ($Im\omega > 0$) provided the normalized Debye length ($\Gamma_D = \lambda_{D\perp}/r_b$) is sufficiently large and the normalized beam intensity ($s_b = \hat{\omega}_{pb}^2/2\gamma_b^2\omega_{\beta\perp}^2$) is sufficiently below the space-charge limit. Depending on system parameters, the growth rate can be a substantial fraction of the focusing frequency $\omega_{\beta\perp}$. As a general remark, application of a warm-fluid model to describe the equilibrium and stability properties of intense charged particle beams appears to be a remarkably robust and simple approach, both for the case of stable high-frequency collective oscillations²⁴, as well as the unstable case considered here, where the instability is driven by gross macroscopic properties of the beam equilibrium (pressure anisotropy).

ACKNOWLEDGMENTS

This research was supported by the U.S. Department of Energy.

REFERENCES

- ¹R. C. Davidson, *Physics of Nonneutral Plasmas* (Addison-Wesley, Reading, MA, 1990), and references therein.
- ² T. P. Wangler, Principles of RF Linear Accelerators (Wiley, New York, 1998).
- ³ M. Reiser, Theory and Design of Charged Particle Beams (Wiley, New York, 1994).
- ⁴I. Kapchinskij and V. Vladimirskij, in Proceedings of the International Conference on High Energy Accelerators and Instrumentation (CERN Scientific Information Service, Geneva, 1959), p. 274.
- ⁵ R. L. Gluckstern, in Proceedings of the 1970 Proton Linear Accelerator Conference, Batavia, IL, 1971, edited by M. R. Tracy (National Accelerator Laboratory, Batavia, IL, 1971), p. 811.
- ⁶T.-S. Wang and L. Smith, Part. Accel. **12**, 247 (1982).
- ⁷I. Hofmann, L. J. Laslett, L. Smith and I. Haber, Part. Accel. **13**, 145 (1983).
- ⁸I. Hofmann and J. Struckmeier, Part. Accel. **21**, 69 (1987).
- ⁹R. L. Gluckstern, W. -H. Cheng and H. Yee, Phys. Rev. Lett. **75**, 2835 (1995).
- ¹⁰C. Chen, R. Pakter and R. C. Davidson, Phys. Rev. Lett. **79**, 225 (1997).
- ¹¹ J. Struckmeier and I. Hofmann, Part. Accel. **39**, 219 (1992).
- ¹² N. Brown and M. Reiser, Phys. Plasmas **2**, 965 (1995).
- ¹³ R. C. Davidson and C. Chen, Part. Accel. **59**, 175 (1998).
- ¹⁴ R. C. Davidson, Phys. Rev. Lett. **81**, 991 (1998).
- ¹⁵ R. C. Davidson, H. Qin and P. J. Channell, Phys. Rev. Special Topics on Accelerators and Beams 2, 074401 (1999).

- ¹⁶ R. C. Davidson and H. Qin, Phys. Rev. Special Topics on Accelerators and Beams 2, 114401 (1999).
- 17 See, e.g., Chaps. 2, 4, 9 and 10 of Ref. 1.
- ¹⁸S. M. Lund, D. A. Callahan, A. Friedman, D. P. Grote, I. Haber and T. -S. Wang, Proceedings of the 19th International Linac Conference (Chicago, IL, 1998).
- ¹⁹I. Haber, D. A. Callahan, A. Friedman, D. P. Grote, S. M. Lund and T. -S. Wang, Nucl. Instrum. Methods Phys. Res. (in press).
- ²⁰I. Haber, D. A. Callahan, A. Friedman, D. P. Grote and A. B. Langdon, J. Fusion Eng. Design **32**, 159 (1996).
- ²¹ Reference 1 presents a general derivation of the macroscopic fluid Maxwell equations from the Vlasov-Maxwell equations on pages 22-26. Several aspects of cold-fluid equilibrium and stability properties of nonneutral beam-plasma systems are described on pages 240-276 of Ref 1.
- ²² R. C. Davidson, P. Stoltz and C. Chen, Phys. Plasmas 4, 3710 (1997).
- ²³I. Hofmann, Phys. Rev. **E57**, 4713 (1998).
- ²⁴S. M. Lund and R. C. Davidson, Phys. Plasmas 5, 3028 (1998).
- ²⁵ C. Chen and R. Pakter, Phys. Plasmas 7, in press (2000).
- ²⁶ R. C. Davidson and S. Strasburg, "Warm-Fluid Stability Properties of Intense Nonneutral Charged Particle Beams with Pressure Anisotropy," submitted for publication (2000).

FIGURES

FIG. 1. Plot of $n^0(r)/\hat{n}$ versus r/r_b calculated from Eqs. (9) and (10) for several values of $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2\omega_{\beta\perp}^2$.

FIG. 2. Plots of (a) $Re\omega_n/\nu_0$ versus ν/ν_0 for radial mode numbers $n = 1, \dots 4$, and (b) $\delta \hat{E}_r = -(\partial/\partial r)\delta \hat{\phi}$ versus r/r_b for n = 1, 2, 3, for $s_b = 0.36$ and $\nu/\nu_0 = 0.8$, obtained numerically from the eigenvalue equation (15) for $k_z = 0$.

FIG. 3. Plots of (a) $Re(\omega - k_z V_b)/\nu_0$ versus $k_z r_b$, and (b) $Re[\delta \hat{\phi}(r)]$ versus r/r_b for several values of $k_z r_b$, obtained numerically from Eq. (15) for $\Gamma_D = 0.360$ ($s_b = 0.755$).

FIG. 4. Plots of (a) $Re(\omega - k_z V_b)/\nu_0$ and $Im\omega/\nu_0$ versus $k_z r_b$, and (b) $Re[\delta \hat{\phi}(r)]$ and $Im[\delta \phi(\hat{r})]$ versus r/r_b for $k_z r_b = 4$, obtained numerically from Eq. (15) for $\Gamma_D = 0.509$ ($s_b = 0.55$).

FIG. 5. Plots of $Im\omega/\nu_0$ versus $k_z r_b$ obtained numerically from Eq. (15) for several values of $\Gamma_D > \Gamma_D^* = 0.364$.



FIG. 1. Plot of $n^0(r)/\hat{n}$ versus r/r_b calculated from Eqs. (9) and (10) for several values of $s_b = \hat{\omega}_{pb}^2/2\gamma_b^2\omega_{\beta\perp}^2$.



FIG. 2. Plots of (a) $Re\omega_n/\nu_0$ versus ν/ν_0 for radial mode numbers $n = 1, \dots 4$, and (b) $\delta \hat{E}_r = -(\partial/\partial r)\delta \hat{\phi}$ versus r/r_b for n = 1, 2, 3, for $s_b = 0.36$ and $\nu/\nu_0 = 0.8$, obtained numerically from the eigenvalue equation (15) for $k_z = 0$.



FIG. 3. Plots of (a) $Re(\omega - k_z V_b)/\nu_0$ versus $k_z r_b$, and (b) $Re[\delta \hat{\phi}(r)]$ versus r/r_b for several values of $k_z r_b$, obtained numerically from Eq. (15) for $\Gamma_D = 0.360$ ($s_b = 0.755$).



FIG. 4. Plots of (a) $Re(\omega - k_z V_b)/\nu_0$ and $Im\omega/\nu_0$ versus $k_z r_b$, and (b) $Re[\delta\hat{\phi}(r)]$ and $Im[\delta\phi(\hat{r})]$ versus r/r_b for $k_z r_b = 4$, obtained numerically from Eq. (15) for $\Gamma_D = 0.509$ ($s_b = 0.55$).



FIG. 5. Plots of $Im\omega/\nu_0$ versus $k_z r_b$ obtained numerically from Eq. (15) for several values of $\Gamma_D > \Gamma_D^* = 0.364$.

The Princeton Plasma Physics Laboratory is operated by Princeton University under contract with the U.S. Department of Energy.

> Information Services Princeton Plasma Physics Laboratory P.O. Box 451 Princeton, NJ 08543

Phone: 609-243-2750 Fax: 609-243-2751 e-mail: pppl_info@pppl.gov Internet Address: http://www.pppl.gov