

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,  
UNDER CONTRACT DE-AC02-76CH03073

PPPL-3980  
UC-70



PPPL-3980

**Physics Results from the  
National Spherical Torus Experiment**

by

M.G. Bell for the NSTX Research Team

July 2004



**PRINCETON PLASMA PHYSICS LABORATORY  
PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY**

## **PPPL Reports Disclaimer**

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

## **Availability**

This report is posted on the U.S. Department of Energy's Princeton Plasma Physics Laboratory Publications and Reports web site in Fiscal Year 2004. The home page for PPPL Reports and Publications is: [http://www.pppl.gov/pub\\_report/](http://www.pppl.gov/pub_report/)

DOE and DOE Contractors can obtain copies of this report from:

U.S. Department of Energy  
Office of Scientific and Technical Information  
DOE Technical Information Services (DTIS)  
P.O. Box 62  
Oak Ridge, TN 37831

Telephone: (865) 576-8401

Fax: (865) 576-5728

Email: [reports@adonis.osti.gov](mailto:reports@adonis.osti.gov)

This report is available to the general public from:

National Technical Information Service  
U.S. Department of Commerce  
5285 Port Royal Road  
Springfield, VA 22161

Telephone: 1-800-553-6847 or  
(703) 605-6000

Fax: (703) 321-8547

Internet: <http://www.ntis.gov/ordering.htm>

# Physics Results from the National Spherical Torus Experiment

M.G. Bell for the NSTX Research Team

Princeton Plasma Physics Laboratory, Princeton, NJ, U.S.A.

The National Spherical Torus Experiment (NSTX) produces plasmas with aspect ratio  $A \equiv R/a = 0.85\text{m}/0.68\text{m} \sim 1.25$ , at plasma currents up to 1.5 MA with vacuum toroidal magnetic field up to 0.6 T on axis. The plasmas are heated by up to 6 MW of High-Harmonic Fast Waves (HHFW) at a frequency 30 MHz and by 7 MW of deuterium Neutral Beam Injection (NBI) at an energy up to 100 keV. Since January 2004, NSTX has been operating, routinely at toroidal fields up to 0.45 T, with a new central conductor bundle in the toroidal field coil.

Recent experiments have concentrated on optimizing and understanding the beta limits through shaping the plasma cross-section and controlling the plasma profiles. The maximum elongation of the plasma cross section has been increased to  $\kappa = 2.6$  and the triangularity (average of upper and lower) to  $\delta = 0.8$ . This was achieved by improving the feedback control for vertical stability [1] and by employing techniques to broaden the plasma current profile. These techniques include early application of the NBI heating and inducing an early transition to the H-mode by a brief pause in the plasma current ramp. The lower plasma inductance resulting from the high  $\kappa$  and broader current profile has the additional benefit of

allowing longer, higher-current pulses within the constraint of the flux swing available from the central solenoid. Toroidal beta up to  $\beta_{T,\text{EFIT}} = \langle p_{\parallel} + p_{\perp} \rangle / 2 / (B_{T0}^2 / 2\mu_0) = 40\%$  as determined by analysis of the equilibrium with EFIT based external magnetic measurements with the pressure profile shape constrained by the electron pressure  $p_e(R)$  measured by Thomson scattering [2]. Figure 1 shows the achieved  $\beta_T$  plotted against normalized plasma current; values of the normalized beta  $\beta_N = \beta_T / (I_p / aB_T) = 6.2 \text{ \%} \cdot \text{m} \cdot \text{T} / \text{MA}$  have been reached in these low-aspect-ratio plasmas. The gains in normalized current from the higher  $\kappa$  and  $\delta$  achieved in recent operation are evident. Analysis of the profile data with the TRANSP code [3] yields  $\beta_{T,\text{TRANSP}} = \langle p_{\parallel} + 2p_{\perp} \rangle / 3 / (B_{T0}^2 / 2\mu_0) = 34\%$  using a classical Monte-Carlo model for the thermalization of the energetic ions from NBI. The waveforms of the shot with highest  $\beta_T$  are shown in Fig. 2 and the profiles of  $T_i$ ,  $T_e$ ,  $n_e$  and  $v_i$ , the toroidal ion velocity, and a cross-section of the plasma equilibrium at the time of maximum  $\beta_T$  are shown in Fig. 3. Note that the very high ion toroidal rotation velocity relative to the Alfvén velocity,  $v_i/v_A \sim 0.3$ , affects the




Fig. 1 Toroidal beta from EFIT vs. Troyon-normalized plasma current.




Fig. 2 Waveforms for the NSTX shot achieving the highest  $\beta_T$ .

equilibrium significantly, leading to an outward shift of the density profile with respect to the flux surfaces [4]. This lower-single-null divertor discharge was initiated at higher toroidal magnetic field to achieve the most quiescent current ramp phase before reducing the field to achieve the highest  $\beta_T$ . The mechanisms responsible for the beta limit in different operating conditions are discussed in [5] and the effects of MHD on fast-ion confinement in [6].




Fig. 3 Profiles of the ion and electron temperature and ion toroidal velocity and electron density at peak  $\beta_T$ .

Boronization is applied in NSTX, at intervals of 1 day to 2 weeks, to control impurities [7] and helium glow discharge cleaning is performed between plasma shots to remove deuterium from plasma-facing surfaces. In addition, prior to the highest  $\beta$  shot, an ohmically heated helium discharge had been run to reduce the recycling further and control the edge density which otherwise can rise rapidly following the H-mode transition. To improve the gas fueling efficiency, a supersonic gas injection nozzle, capable of injecting a collimated jet at  $\sim 1.5$  km/s at the outside midplane is being installed on NSTX [8].

The basic dependences of the total and thermal confinement time in NSTX on plasma current and NBI heating power are similar to those of conventional tokamaks, both in controlled single-parameter scans and in regression fits to the NSTX database. This is shown in Fig. 4, where a dependence of the total confinement time from the EFIT magnetic analysis,  $\tau_E \propto I_p^{0.84 \pm 0.05} P_{tot}^{-0.49 \pm 0.04}$  is inferred for H-mode discharges spanning the ranges  $0.6 \leq I_p [\text{MA}] \leq 1.2$ ,  $1.5 \leq P_{NB} [\text{MW}] \leq 7$ . However, the total confinement times exceed the ITER-97L [9] scaling for confinement by factors 1.5 – 2.5 for discharges both with and without H-mode transitions. It is particularly encouraging that enhancements of the




Fig. 4 Fits to the magnetically determined energy confinement time for H-mode deuterium discharges with  $B_T = 0.45$  T,  $\kappa = 1.8 - 2.4$  at the time of peak total electron energy.

a)  $\tau_E \propto I_p^{+0.78}$  at constant  $P_{tot}$ , b)  $\tau_E \propto P_{tot}^{-0.47}$  at constant  $I_p$ , c) Regression to both  $I_p$ ,  $P_{tot}$ .

confinement over ITER-97L scaling by a factor  $H_{97L} = 2$  have been maintained as the density was raised by gas puffing to the Greenwald limit. The thermal confinement times have been analyzed by TRANSP for a smaller dataset to date, but in H-mode plasmas these too exceed the corresponding ITER-98Hpb(y,2) scaling.

The TRANSP analysis shows that the dominant thermal loss is through the electron channel in both H- and L- mode plasmas. This situation is consistent with gyro-kinetic calculations of the plasma stability [10] which show that the ion temperature gradient modes may be stabilized by the large shear in the toroidal rotation. Figure 5 shows the measured profiles and the thermal diffusivities for the ions and electrons calculated by TRANSP for a low-density L-mode discharge which shows evidence for an internal transport barrier, *i.e.* a region of reduced  $\chi_e$  and  $\chi_i$ , near  $r/a \approx 0.5$ . Although this shot did not show an H-mode transition, its confinement was enhanced over L-mode by a factor  $H_{97L} \approx 2.4$ . The relationship between this transport reduction and possible reversal of the magnetic shear,




Fig. 5 a) Profiles of the ion and electron temperature, ion toroidal velocity and electron density and b) profiles of the ion and electron and neoclassical ion thermal diffusivities for a low-density L-mode plasma with an apparent internal transport barrier.

will be investigated using the motional-Stark-effect diagnostic which is now being commissioned on NSTX. In some high-density ( $n_e(0) \sim 8 \times 10^{19} m^{-3}$ ) NBI-heated, H-mode plasmas, recent analyses [11] of the power balance taking account of the experimental uncertainties have confirmed that the ion thermal diffusivity drops to the neoclassical level, or even below in some cases.

Significant electron heating has been produced by HHFW power in low-density deuterium and helium plasmas. The most effective heating has been observed when the toroidal array of 12 poloidal coupling straps has been phased to launch waves with a symmetrical toroidal wavenumber spectrum peaked at  $k_{tor} \approx \pm 14 m^{-1}$  or  $\pm 7 m^{-1}$ . The HHFW heating can induce H-mode transitions in deuterium plasmas and, as with NBI, there is a striking buildup of the density in the edge following the H-mode transition. However, the overall confinement in the plasmas heated by HHFW is lower than for NBI heating. Figure 6 shows the measured plasma total and the electron energy and the prediction based on ITER-97L scaling assuming that all the launched RF power is coupled to the




Fig. 6 Measured total and electron energies in deuterium plasmas heated by HHFW.

plasma. The H-mode cases tend to exceed the L-mode plasmas in total and electron energy but the central electron temperatures in the H-mode actually tend to be lower, suggesting that less power is deposited in the center with H-mode profiles. The HHFW frequency is about 9 times the deuteron cyclotron frequency at the plasma axis, so little direct absorption of the waves by the thermal ions is expected in NSTX, although an interaction of the HHFW with the energetic ions from NBI has been previously observed and explained [12]. Experiments are underway to determine whether the difference in the effect on the plasma between the HHFW and the NBI heating is caused by the difference in the division of the heating power between the electrons and ions, or whether there are additional absorption or loss mechanisms at work with HHFW heating. A spectrometer measuring helium emission from the plasma edge has revealed the development of a substantial hot ion population, with perpendicular temperatures up to  $\sim 0.5$  keV, during HHFW heating [13]. Evidence has also been obtained from a Langmuir probe with high frequency response inserted into the far scrape-off layer, for a possible parametric decay of the launched waves into ion Bernstein waves at harmonics of the local deuteron cyclotron frequency. When the HHFW coupler array was phased to produce co- and counter- directed waves with  $k_{\text{tor}} \approx \pm (3.5 - 7) \text{ m}^{-1}$  in pairs of otherwise matched plasmas, differences were measured in the toroidal loop voltage (corrected for inductive effects), indicating current drive estimated to be about 100 kA [14].

In H-mode plasmas, a variety of ELM activity occurs; examples are seen in the  $D_\alpha$  light trace in Fig. 2. After a quiescent phase following the H-mode transition, small frequent ELMs occur between 0.25 and 0.35 s. These ELMs have characteristics different from other well-known types and have tentatively been designated as “Type V” [15]. The large (Type I) ELMs which begin at about 0.35 s become more frequent as the toroidal field is reduced, whereas in a companion shot with constant toroidal field, they remained sporadic.

The presenting author wishes to thank colleagues in the NSTX research team for providing data and for helpful discussion of these results. This work is supported by U.S. DOE Contract DE-AC02-76CH03073.

## References

- [1] D. Mueller *et al.*, paper P2-199, *this conference*.
- [2] S.A. Sabbagh *et al.*, Phys. Plasmas **9** (2002) 2085
- [3] J. Ongena *et al.*, Fusion Technology **33** (1997) 181
- [4] J. Menard, Nucl. Fusion **43** (2003) 330.
- [5] A.C. Sontag *et al.*, paper P2-193, *this conference*.
- [6] S.S. Medley *et al.*, paper P2-200, *this conference*.
- [7] H.W. Kugel *et al.*, J. Nucl. Mater. **313–316** (2003) 187
- [8] V. Soukhanovskii *et al.*, paper P2-190, *this conference*.
- [9] ITER Physics Expert Groups, Nuclear Fusion **39** (1999) 2175
- [10] M. Redi *et al.*, paper P2-162, *this conference*.
- [11] B.P. LeBlanc *et al.* Nuclear Fusion **44** (2004) 513.
- [12] A.L. Rosenberg *et al.*, Physics of Plasmas, **11** (2004) 2441-2452.
- [13] T.M. Biewer *et al.*, paper P2-198, *this conference*.
- [14] J.R. Wilson, Phys. Plasmas **10** (2003) 1733
- [15] R. Maingi *et al.*, submitted to J. Nucl. Mater., May 2004 and paper P2-189, *this conference*.

## External Distribution

Plasma Research Laboratory, Australian National University, Australia  
Professor I.R. Jones, Flinders University, Australia  
Professor João Canalle, Instituto de Fisica DEQ/IF - UERJ, Brazil  
Mr. Gerson O. Ludwig, Instituto Nacional de Pesquisas, Brazil  
Dr. P.H. Sakanaka, Instituto Fisica, Brazil  
The Librarian, Culham Laboratory, England  
Mrs. S.A. Hutchinson, JET Library, England  
Professor M.N. Bussac, Ecole Polytechnique, France  
Librarian, Max-Planck-Institut für Plasmaphysik, Germany  
Jolan Moldvai, Reports Library, Hungarian Academy of Sciences, Central Research Institute  
for Physics, Hungary  
Dr. P. Kaw, Institute for Plasma Research, India  
Ms. P.J. Pathak, Librarian, Institute for Plasma Research, India  
Ms. Clelia De Palo, Associazione EURATOM-ENEA, Italy  
Dr. G. Grosso, Instituto di Fisica del Plasma, Italy  
Librarian, Naka Fusion Research Establishment, JAERI, Japan  
Library, Laboratory for Complex Energy Processes, Institute for Advanced Study,  
Kyoto University, Japan  
Research Information Center, National Institute for Fusion Science, Japan  
Dr. O. Mitarai, Kyushu Tokai University, Japan  
Dr. Jiengang Li, Institute of Plasma Physics, Chinese Academy of Sciences,  
People's Republic of China  
Professor Yuping Huo, School of Physical Science and Technology, People's Republic of China  
Library, Academia Sinica, Institute of Plasma Physics, People's Republic of China  
Librarian, Institute of Physics, Chinese Academy of Sciences, People's Republic of China  
Dr. S. Mirnov, TRINITI, Troitsk, Russian Federation, Russia  
Dr. V.S. Strelkov, Kurchatov Institute, Russian Federation, Russia  
Professor Peter Lukac, Katedra Fyziky Plazmy MFF UK, Mlynska dolina F-2,  
Komenskeho Univerzita, SK-842 15 Bratislava, Slovakia  
Dr. G.S. Lee, Korea Basic Science Institute, South Korea  
Institute for Plasma Research, University of Maryland, USA  
Librarian, Fusion Energy Division, Oak Ridge National Laboratory, USA  
Librarian, Institute of Fusion Studies, University of Texas, USA  
Librarian, Magnetic Fusion Program, Lawrence Livermore National Laboratory, USA  
Library, General Atomics, USA  
Plasma Physics Group, Fusion Energy Research Program, University of California  
at San Diego, USA  
Plasma Physics Library, Columbia University, USA  
Alkesh Punjabi, Center for Fusion Research and Training, Hampton University, USA  
Dr. W.M. Stacey, Fusion Research Center, Georgia Institute of Technology, USA  
Dr. John Willis, U.S. Department of Energy, Office of Fusion Energy Sciences, USA  
Mr. Paul H. Wright, Indianapolis, Indiana, USA

The Princeton Plasma Physics Laboratory is operated  
by Princeton University under contract  
with the U.S. Department of Energy.

Information Services  
Princeton Plasma Physics Laboratory  
P.O. Box 451  
Princeton, NJ 08543

Phone: 609-243-2750  
Fax: 609-243-2751  
e-mail: [pppl\\_info@pppl.gov](mailto:pppl_info@pppl.gov)  
Internet Address: <http://www.pppl.gov>