


PPPL-4083

PPPL-4083

Testing Gyrokinetics on C-Mod and NSTX

M.H. Redi, W. Dorland, C.L. Fiore, D. Stutman, J.A. Baumgaertel,
B. Davis, S.M. Kaye, D.C. McCune, J. Menard, and G. Rewoldt

June 2005



PPPL Report Disclaimers

Full Legal Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Trademark Disclaimer

Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors.

PPPL Report Availability

This report is posted on the U.S. Department of Energy's Princeton Plasma Physics Laboratory Publications and Reports web site in Fiscal Year 2005. The home page for PPPL Reports and Publications is: http://www.pppl.gov/pub_report/

Office of Scientific and Technical Information (OSTI):

Available electronically at: <http://www.osti.gov/bridge>.

Available for a processing fee to U.S. Department of Energy and its contractors, in paper from:

U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831-0062
Telephone: (865) 576-8401
Fax: (865) 576-5728
E-mail: reports@adonis.osti.gov

National Technical Information Service (NTIS):

This report is available for sale to the general public from:

U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
Telephone: (800) 553-6847
Fax: (703) 605-6900
Email: orders@ntis.fedworld.gov
Online ordering: <http://www.ntis.gov/ordering.htm>

Testing Gyrokinetics on C-Mod and NSTX

M. H. Redi¹, W. Dorland², C. L. Fiore³, D. Stutman⁴, J. A. Baumgaertel¹, B. Davis¹,
S. M. Kaye¹, D. C. McCune¹, J. Menard¹, G. Rewoldt¹

¹Princeton Plasma Physics Laboratory, Princeton University, Princeton, NJ 08540 USA

²University of Maryland, College Park, MD 20742 USA

³Plasma Science Fusion Center, MIT, Cambridge, MA 02139 USA

⁴Johns Hopkins University, Baltimore, MD 21218 USA

Quantitative benchmarks of computational physics codes against experiment are essential for the credible application of such codes. Fluctuation measurements can provide necessary critical tests of nonlinear gyrokinetic simulations, but such require extraordinary computational resources. Linear microstability calculations with the GS2 [1] gyrokinetic code have been carried out for tokamak and ST experiments which exhibit internal transport barriers (ITB) and good plasma confinement. Qualitative correlation is found for improved confinement before and during ITB plasmas on Alcator C-Mod [2] and NSTX [3] with weaker long wavelength microinstabilities in the plasma core regions. Mixing length transport models are discussed. The NSTX L-mode is found to be near marginal stability for kinetic ballooning modes.

Fully electromagnetic, linear, gyrokinetic calculations of the Alcator C-Mod ITB during off-axis rf heating, following four plasma species and including the complete electron response show ITG/TEM microturbulence is suppressed in the plasma core and in the barrier region before barrier formation, without recourse to the usual requirements of velocity shear or reversed magnetic shear [4-5]. No strongly growing long or short wavelength drift modes are found in the plasma core but strong ITG/TEM and ETG drift wave turbulence is found outside the barrier region. Linear microstability analysis is qualitatively consistent with the experimental transport analysis, showing low transport inside and high transport outside the ITB region before barrier formation, without consideration of \mathbf{ExB} shear stabilization.

Calculations of the ITG mode instability threshold and the dependence of instability on temperature and density scaling length have been used to test the GS2 gyrokinetic model for ITG microturbulence on C-Mod [4-5]. Transport analysis at the ITB onset time in H-mode experiments [6] is in rough agreement with the linear mixing length model $\gamma < k_{\perp}^2$. The mixing length estimates exceed experimental estimates of transport by less than a factor of two and are in agreement with experiment in the plasma core (see Table I). GS2

calculations were also used to verify an algebraic, computationally-based model for the ITG instability threshold [7] and standard analytic models [8, 9]. Agreement of these threshold models (within 40%) with the critical temperature threshold obtained with the more complete physics of the GS2 code verifies that these standard models are approximately correct. Figure 1 shows a phase diagram for ITG/TEM instability for C-Mod at ITB onset in the barrier region on which are mapped the locations for linear GS2 calculations. The topology of the diagram, adapted from Ref. 10, is preserved but for this case, the critical ITG temperature threshold is 6.9, rather than 2.5 as in Ref. 10. Considering experimental errors of 10-20%, and even larger errors in plasma gradients, these differences are not remarkable.

Definitive testing of gyrokinetics is much more difficult for NSTX than for C-Mod, since the potentially most dangerous, long wavelength modes are not ITG/TEM only. The new microtearing modes may cause significant transport. NSTX L-mode plasmas are found to exhibit improved electron transport [11] and less strongly destabilized microtearing than do H-modes. ETG and microtearing modes dominate in different regions of NSTX plasmas and so appear to compete for the same free energy source. Further, new calculations show that L-mode plasma is just below the critical threshold for kinetic ballooning modes (Fig. 2). The linear microstability of a slow current ramp, low density L-mode pulse #112996 was studied at a selected radius, $r/a=0.55$, using an equilibrium based on experimental data. At this radius, where the local beta is 9%, the ITG/TEM modes are unstable, and the KBM is near marginal stability. A self-consistent beta scan, based on a Miller model MHD equilibrium, indicates that the critical beta is 11% for strongly growing kinetic ballooning modes. The toroidal mode numbers and the real frequency are of the order of those expected for the KBM, with the growth rates peaking when the mode number is in the range 6 to 9.

Initial gyrokinetic calculations of this NSTX L-mode at an earlier time [11] show that for $r/a = 0.35$ the microtearing is maximum at $0.3 k_{\perp}\rho_s$. At $r/a = 0.45$ the maximum ITG growth rate occurs at $0.3 k_{\perp}\rho_s$ and ETG peaks at $20 k_{\perp}\rho_s$. At $r/a = 0.65$, ITG is most strongly growing at $0.7 k_{\perp}\rho_s$ and ETG is maximum at $30 k_{\perp}\rho_s$. Other modes are not found to be unstable.

The effect of temperature gradients and magnetic shear variations on microstability was also studied at $r/a=0.35$ with fully electromagnetic calculations following four plasma species and including the complete electron response. In this case however, analytic Miller-type equilibria were used, rather than numerical equilibria. At this radius the local β is 19%. The calculations lead to microtearing modes and ITG/TEM modes, in addition to other

electromagnetic modes of longer wavelength and even parity, which rotate in the ion diamagnetic direction (possibly kink modes). Core microtearing modes have been found unstable in every NSTX plasma examined, unless the plasma has reversed shear or is assumed collisionless [12, 13]. Figures 3-4 show the effects of a scan of both magnetic shear and temperature scale length variations on the ITG/TEM and microtearing long wavelength modes. As expected, microtearing is destabilized for positive temperature gradients and low magnetic shear (Fig. 3) while ITG is destabilized by reversed temperature gradients and low magnetic shear (Fig. 4). A more complex mixing length model than applied to the C-Mod case will be needed for NSTX, to properly include all the long wavelength modes.

*Supported by USDoE contract No. DE-AC02-76CH03073

- [1] M. Kotschenreuther, *et al.*, Comput. Phys. Commun. **88**, 128 (1995).
- [2] I. H. Hutchinson, *et al.*, Phys. Plas. **1**, 1511 (1994).
- [3] M. Ono, *et al.* Proc. 18th Int. Conf. on Fusion Energy (Sorrento) IAEA, Vienna, 2000.
- [4] M. H. Redi *et al.* “Microturbulent drift mode stability before internal transport barrier formation in the Alcator C-Mod radio frequency heated H-mode”, Phys. Plas. in press.
- [5] J. A. Baumgaertel, *et al.*, “Marginal stability of microturbulence near ITB onset on Alcator C-Mod”, DoE OFES Journal of Undergraduate Research, Vol 5 (2005).
- [6] C. L. Fiore, *et al.* Plasma Physics and Controlled Fusion **46**, B281-291 (2004).
- [7] F. Jenko, W. Dorland, G. W. Hammett, Phys. Plas. **8**, 4096 (2001).
- [8] F. Romanelli, Phys. Fluids **B1** 1018 (1989)
- [9] T. S. Hahm, W. M. Tang, Phys. Fluids **B1**, 1185 (1989).
- [10] X. Garbet, P. Mantica, C. Angioni, *et al.*, Plasma Physics and Controlled Fusion, **46** B557-B574 (2004).
- [11] D. Stutman, *et al.*, “Studies of Improved Electron Confinement on NSTX”, submitted to Nucl. Fus. (2004).
- [12] M. H. Redi, *et al.* Proc. 30st EPS Conference on Plasma Physics, St. Petersburg, 2003.
- [13] M. H. Redi, *et al.* Proc. 31st EPS Conference on Plasma Physics, London, 2004.

Table I. Comparison of linear ITG mixing length model transport coefficients to anomalous transport coefficients derived from transport analysis of experiment

	r/a	0.25	0.45	0.65
$\chi_{mix} = \gamma / \langle k_{\perp}^2 \rangle$ (m^2/s)		0	0.76	1.04
$\chi_{eff} - \chi_i^{Chang-Hinton}$ (m^2/s)		0	0.4	0.8




Fig. 1. Phase diagram for linear instability of ITG and TEM drift modes for the C-Mod plasma before ITB formation. Diagram adapted from Fig. 1 of Ref. 10. Star symbols indicate three C-Mod radial locations simulated, with stable conditions found in the plasma core (red), weakly unstable ITG at the barrier region (green) and strongly unstable ITG outside the barrier region (blue). Dashed lines map locations for linear simulations carried out for variations of the barrier region.




Fig. 2 Growth rates and real frequencies of fastest growing mode in NSTX shot 112996 at 0.4 sec, $r/a=0.55$. A beta scan shows that the experiment is dominated by ITG/TEM and is just below the critical threshold for the kinetic ballooning mode. The KBM may be limiting beta at this radius, $\beta_{crit} \sim 11\%$.




Fig. 3 Growth rates for microtearing mode identified in calculations of NSTX L-mode plasmas with variations in magnetic shear and scaled temperature gradients of all four species. Mode is destabilized at low magnetic shear and positive temperature gradients.




Fig. 4 Growth rates for ion temperature gradient modes identified in calculations of NSTX L-mode plasmas for variations in magnetic shear and scaled temperature gradients of all four species. Mode is destabilized at low magnetic shear for increased temperature gradients of positive and negative sign.

External Distribution

Plasma Research Laboratory, Australian National University, Australia
Professor I.R. Jones, Flinders University, Australia
Professor João Canalle, Instituto de Fisica DEQ/IF - UERJ, Brazil
Mr. Gerson O. Ludwig, Instituto Nacional de Pesquisas, Brazil
Dr. P.H. Sakanaka, Instituto Fisica, Brazil
The Librarian, Culham Science Center, England
Mrs. S.A. Hutchinson, JET Library, England
Professor M.N. Bussac, Ecole Polytechnique, France
Librarian, Max-Planck-Institut für Plasmaphysik, Germany
Jolan Moldvai, Reports Library, Hungarian Academy of Sciences, Central Research
Institute for Physics, Hungary
Dr. P. Kaw, Institute for Plasma Research, India
Ms. P.J. Pathak, Librarian, Institute for Plasma Research, India
Dr. Pandji Triadyaksa, Fakultas MIPA Universitas Diponegoro, Indonesia
Professor Sami Cuperman, Plasma Physics Group, Tel Aviv University, Israel
Ms. Clelia De Palo, Associazione EURATOM-ENEA, Italy
Dr. G. Grosso, Instituto di Fisica del Plasma, Italy
Librarian, Naka Fusion Research Establishment, JAERI, Japan
Library, Laboratory for Complex Energy Processes, Institute for Advanced Study,
Kyoto University, Japan
Research Information Center, National Institute for Fusion Science, Japan
Professor Toshitaka Idehara, Director, Research Center for Development of Far-Infrared Region,
Fukui University, Japan
Dr. O. Mitarai, Kyushu Tokai University, Japan
Mr. Adefila Olumide, Ilorin, Kwara State, Nigeria
Dr. Jiangang Li, Institute of Plasma Physics, Chinese Academy of Sciences, People's Republic of China
Professor Yuping Huo, School of Physical Science and Technology, People's Republic of China
Library, Academia Sinica, Institute of Plasma Physics, People's Republic of China
Librarian, Institute of Physics, Chinese Academy of Sciences, People's Republic of China
Dr. S. Mirnov, TRINITI, Troitsk, Russian Federation, Russia
Dr. V.S. Strelkov, Kurchatov Institute, Russian Federation, Russia
Kazi Firoz, UPJS, Kosice, Slovakia
Professor Peter Lukac, Katedra Fyziky Plazmy MFF UK, Mlynska dolina F-2, Komenskeho Univerzita,
SK-842 15 Bratislava, Slovakia
Dr. G.S. Lee, Korea Basic Science Institute, South Korea
Dr. Rasulkhozha S. Sharafiddinov, Theoretical Physics Division, Institute of Nuclear Physics, Uzbekistan
Institute for Plasma Research, University of Maryland, USA
Librarian, Fusion Energy Division, Oak Ridge National Laboratory, USA
Librarian, Institute of Fusion Studies, University of Texas, USA
Librarian, Magnetic Fusion Program, Lawrence Livermore National Laboratory, USA
Library, General Atomics, USA
Plasma Physics Group, Fusion Energy Research Program, University of California at San Diego, USA
Plasma Physics Library, Columbia University, USA
Alkesh Punjabi, Center for Fusion Research and Training, Hampton University, USA
Dr. W.M. Stacey, Fusion Research Center, Georgia Institute of Technology, USA
Director, Research Division, OFES, Washington, D.C. 20585-1290

The Princeton Plasma Physics Laboratory is operated
by Princeton University under contract
with the U.S. Department of Energy.

Information Services
Princeton Plasma Physics Laboratory
P.O. Box 451
Princeton, NJ 08543

Phone: 609-243-2750
Fax: 609-243-2751
e-mail: pppl_info@pppl.gov
Internet Address: <http://www.pppl.gov>