

PREPARED FOR THE U.S. DEPARTMENT OF ENERGY,
UNDER CONTRACT DE-AC02-76CH03073

PPPL-3524
UC-70



PPPL-3524

Strong "Quantum" Chaos in the Global Ballooning Mode
Spectrum of Three-dimensional Plasmas

by

R.L. Dewar, P. Cuthbert, and R. Ball

November 2000



PRINCETON PLASMA PHYSICS LABORATORY
PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY

PPPL Reports Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Availability

This report is posted on the U.S. Department of Energy's Princeton Plasma Physics Laboratory Publications and Reports web site in Calendar Year 2000. The home page for PPPL Reports and Publications is: http://www.pppl.gov/pub_report/

DOE and DOE Contractors can obtain copies of this report from:

U.S. Department of Energy
Office of Scientific and Technical Information
DOE Technical Information Services (DTIS)
P.O. Box 62
Oak Ridge, TN 37831

Telephone: (865) 576-8401
Fax: (865) 576-5728
Email: reports@adonis.osti.gov

This report is available to the general public from:

National Technical Information Service
U.S. Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

Telephone: 1-800-553-6847 or
(703) 605-6000
Fax: (703) 321-8547
Internet: <http://www.ntis.gov/ordering.htm>

Strong “quantum” chaos in the global ballooning mode spectrum of three-dimensional plasmas

R. L. Dewar*

Princeton University Plasma Physics Laboratory, P.O. Box 451, Princeton N.J. 08543

P. Cuthbert and R. Ball

Department of Theoretical Physics and Plasma Research Laboratory, Research School of Physical Sciences & Engineering, The Australian National University, Canberra 0200 Australia

(November 27, 2000)

The spectrum of ideal magnetohydrodynamic (MHD) pressure-driven (ballooning) modes in strongly nonaxisymmetric toroidal systems is difficult to analyze numerically owing to the singular nature of ideal MHD caused by lack of an inherent scale length. In this paper, ideal MHD is regularized by using a k -space cutoff, making the ray tracing for the WKB ballooning formalism a chaotic Hamiltonian billiard problem. The minimum width of the toroidal Fourier spectrum needed for resolving toroidally localized ballooning modes with a global eigenvalue code is estimated from the Weyl formula. This phase-space-volume estimation method is applied to ballooning-unstable plasma equilibria in the H-1NF helical axis stellarator and the Large Helical Device (LHD).

PACS numbers: 52.35.Py, 52.55.Hc, 05.45.Mt

In design studies for new magnetic confinement devices for fusion plasma experiments (e.g. investigations [1,2] leading to the proposed National Compact Stellarator Experiment, NCSX [3]), the maximum pressure that can stably be confined in any proposed magnetic field configuration is routinely estimated by treating the plasma as an ideal magnetohydrodynamic (MHD) fluid. One linearizes about a sequence of equilibrium states with increasing pressure, and studies the spectrum of normal modes (frequency ω) to determine when there is a component with $\text{Im}\omega > 0$, signifying instability.

Even with the simplification obtained by using the ideal MHD model, the computational task of determining the theoretical stability of a three-dimensional (i.e. non-axisymmetric) device, such as NCSX or the four currently operating helical axis stellarators [4], remains a challenging one.

The problem can be posed as a Lagrangian field theory, with the potential term being the energy functional δW [5]. For a static equilibrium, the kinetic energy is quadratic in ω , so that ω^2 is real. Thus instability occurs when $\omega^2 < 0$. There are two main approaches to

analyzing the spectrum—local and global.

In the *local* approach, which is used for analytical simplification, one orders the scale length of variation of the eigenfunction across the magnetic field lines to be short compared with equilibrium scale lengths [6]. Both interchange and ballooning stability can be treated by solving the general *ballooning equations* [7], a system of ordinary differential equations defined on a given magnetic field line.

The *global* (Galerkin) approach is to expand the plasma displacement field in a finite basis set, inserting this ansatz in the Lagrangian to find a matrix eigenvalue representation of the spectral problem. This approach has been implemented for ideal MHD in three-dimensional plasmas in two codes, TERPSICHORE [8] and CAS3D [9].

Although the Galerkin approach is potentially exact, if one could use a complete, infinite basis set, it is in practice computationally challenging due to the large number of basis functions required to resolve localized instabilities. This leads to very large matrices which must be diagonalized by iterative methods. There is a need for analytical insight to determine a suitable truncated basis set and to predict the nature of the spectrum, e.g. whether it is continuous or discrete.

Such insight may be obtained by a *hybrid* local-global approach, in which one uses a Wentzel–Brillouin–Kramers (WKB) representation of the eigenfunction. In the short-wavelength limit, the same analytical simplifications as are obtained in the local approach are found to give a local dispersion relation that can be used to give information on the global spectrum by using ray tracing and semiclassical quantization.

In axisymmetric systems [10] or in cases where helical ripple can be averaged out, giving an adiabatic invariant, [11,12], the ray equations are integrable and hence the spectrum is characterized by “good quantum numbers”.

However, it has been known for many years [7] that the ray-tracing problem in strongly three-dimensional systems is singular because, in the absence of an adiabatic invariant, the phase-space motion is not bounded—the rays escape to infinity in the wavevector sector. Dewar and Glasser [7] argued that this gives rise to a *continuous* unstable spectrum, with correspondingly singular

*Permanent address: Research School of Physical Sciences & Engineering, The Australian National University. E-mail: robert.dewar@anu.edu.au.

generalized eigenfunctions. (A more rigorous treatment involves the concept of the essential spectrum and Weyl sequences [13,14].)

Our proposed regularization of this singularity can be understood using a simple quantum analogy. Consider the one-dimensional time-independent Schrödinger equation $H\psi = E\psi$ in the limit as the mass of the particle goes to infinity. Then the kinetic energy disappears and the Hamiltonian becomes $H = V(x)$, where V is the potential energy, assumed here to be the harmonic oscillator potential, $\frac{1}{2}x^2$ in suitable units. In the usual Hilbert space the energy spectrum is continuous: $E \geq 0$ and the (generalized) eigenfunctions singular: $\psi(x) = \delta(x - x_E) \pm \delta(x + x_E)$, where $V(x_E) \equiv E$.

We now seek a regularization of this problem by restricting ψ to the space of functions with a finite bandwidth in wavenumber k :

$$\psi(x) = \int_{-k_{\max}}^{k_{\max}} \frac{dk}{2\pi} \psi_k \exp ikx. \quad (1)$$

This truncated Fourier representation models what occurs when one seeks to find the spectrum numerically using a truncated Fourier representation.

We take as starting point a Lagrangian for the wavefunction,

$$L = \int_{-\infty}^{\infty} \psi^* [E - V(x)] \psi dx. \quad (2)$$

Inserting Eq. (1) in Eq. (2) gives

$$L = \int_{-k_{\max}-0}^{k_{\max}+0} \left[E |\psi_k|^2 - \left| \frac{d\psi_k}{dk} + \psi_k \delta(k + k_{\max}) - \psi_k \delta(k - k_{\max}) \right|^2 \right] \frac{dk}{2\pi}. \quad (3)$$

This is infinite unless we require the coefficients of the δ -functions to vanish. That is, $\psi_k = 0$ at $k = \pm k_{\max}$. The Euler-Lagrange equation is $(d^2/2dk^2 + E)\psi_k = 0$, which has the solutions $\exp \pm i(2E)^{1/2}k$. These waves would propagate to infinity if it were not for the reflecting boundary conditions at $\pm k_{\max}$ we have just derived.

That is, we have removed the continuum by box quantization in k -space. In the following we shall do the same for the ballooning mode problem.

As in [7] we write the magnetic field of an arbitrary three-dimensional toroidal equilibrium plasma with nested magnetic flux surfaces labeled by an arbitrary parameter s as $\mathbf{B} = \nabla\zeta \times \nabla\psi - q\nabla\theta \times \nabla\psi \equiv \nabla\alpha \times \nabla\psi$, where $\alpha \equiv \zeta - q\theta$. Here, θ and ζ are the poloidal and toroidal angles, respectively, $\psi(s)$ is the poloidal flux function, and $q(s)$ is the inverse of the rotational transform. Since $\mathbf{B} \cdot \nabla s = \mathbf{B} \cdot \nabla\alpha = 0$, s and α serve to label an individual field line.

We take the stream function [6] to be given by $\varphi = \widehat{\varphi} \exp(iS - i\omega t)$, where $\widehat{\varphi}(\theta|s, \alpha)$ is assumed to vary on

the equilibrium scale. The phase variation is taken to be rapid, so $\mathbf{k} \equiv \nabla S$ is ordered to be large. The frequency ω is ordered $O(1)$, which requires that the wave vector be perpendicular to \mathbf{B} : $\mathbf{k} \cdot \mathbf{B} \equiv 0$. (In this study we consider unstable ideal MHD modes, $\omega^2 < 0$.)

It immediately follows that the eikonal is constant on each field line: $S = S(\alpha, s)$. From the definition of the wave vector, $\mathbf{k} = k_\alpha \nabla\alpha + k_s \nabla s \equiv k_\alpha [\nabla\alpha + \theta_k q'(s) \nabla s]$ where $k_\alpha \equiv \partial S / \partial \alpha$ and $k_s \equiv \partial S / \partial s$. Here the angle-like ballooning parameter θ_k appears naturally as the ratio $k_s / q'(s) k_\alpha$ [10].

The ballooning equation emerges in the large $|\mathbf{k}|$ expansion [7,6] as an ordinary differential equation to be solved on each field line (α, s) with given (k_α, k_s) under the boundary condition $\widehat{\varphi}(\theta) \rightarrow 0$ at infinity to give the eigenvalue $\lambda(\alpha, s, k_\alpha, k_s)$. This constitutes a local dispersion relation $\lambda \equiv \rho\omega^2$ (the mass density ρ being assumed constant everywhere).

The ray equations are the characteristics of the eikonal equation $\lambda(\alpha, s, \partial_\alpha S, \partial_s S) = \rho\omega^2$. These are Hamiltonian equations of motion with α, s the generalized coordinates, k_α, k_s the canonically conjugate momenta, and λ as the Hamiltonian.

In axi- or helically symmetric systems all field lines on a given magnetic surface are equivalent— α is ignorable and k_α is a constant of the motion. In this case the equations are integrable and semiclassical quantization can be used to predict the approximate spectrum of global ballooning instabilities [10]. This technique can sometimes be applied successfully, even in nonsymmetric systems, if there are regions of phase space with a large measure of invariant tori [15,11]. In [11] this was verified using the global eigenvalue code TERPSICHORE [8].

At the other extreme, if the ray orbits are chaotic (but still bounded) then the global spectrum is not regularly structured, but must rather be described statistically by the density of states and the probability distribution of level spacings using the techniques of quantum chaos theory (see e.g. [16,17]).

However, because of the scale invariance of the ideal MHD equations, λ depends only on the direction of \mathbf{k} , *not* on its magnitude: $\lambda = \lambda(\alpha, s, \theta_k)$. This has the consequence that the ray orbits are *unbounded* in phase space, so, strictly speaking, ideal MHD gives rise to a quantum *chaotic scattering* [16,17] problem rather than a straight quantum *chaos* problem. This leads to the continuous spectrum [7] with singular generalized eigenfunctions that cannot really be represented using the simple eikonal ansatz.

On the other hand, the absence of a natural length scale in ideal MHD is a mathematical artifact. Physically, the ion Larmor radius provides a lower cutoff in space, or an upper cutoff in $|\mathbf{k}|$, beyond which ideal MHD ceases to apply. The ballooning equation is also physically regularized by inclusion of diamagnetic drift [18,15].

However, since in general it leads to a complex ray

tracing problem [19], we shall not attempt to model diamagnetic drift stabilization in this paper. Rather, we regularize the ray equations simply by adding a barrier term to the effective ray “Hamiltonian” $H(\alpha, s, k_\alpha, k_s)$,

$$H = \lambda(\alpha, s, k_\alpha, k_s) + U(k_\alpha), \quad (4)$$

where the barrier potential we use is $U(k_\alpha) \equiv K(|k_\alpha| - k_{\max})^2$ for $|k_\alpha| > k_{\max}$ and 0 for $|k_\alpha| < k_{\max}$. In the limit of the constant $K \rightarrow \infty$, this infinite box potential gives the ideal MHD ray equations for $|k_\alpha| < k_{\max}$ and reflecting boundary conditions at $|k_\alpha| = k_{\max}$. Thus we have a two-degree of freedom Hamiltonian billiard problem.

Although overly crude for modeling FLR stabilization, the cutoff at $|k_\alpha| = k_{\max}$ provides a reasonable model for representing the finite spectral bandwidth in the toroidal Fourier mode number (n) representation used in the global eigenvalue codes TERPSICHORE [8] and CAS3D [9].




FIG. 1. The sections $\theta_k = 0$ and $q = 0.893$ of the topologically spherical isosurfaces of the central, (0,0), ballooning mode branch, bounded by the isosurface $\lambda = -6$ (arbitrary units). The darker shades denote higher growth rates, the peak corresponding to $\lambda \approx -8$.

Using ballooning-unstable plasma equilibria calculated for the H-1NF heliac [20,4] using the VMEC code [21], detailed parameter scans have been undertaken for two cases. The first case studied [22] was obtained by increasing the pressure gradient of a marginally stable equilibrium [23] uniformly across the plasma and thus was ballooning unstable at the edge of the plasma. The ray tracing problem for this case would involve consideration of the effect of the plasma boundary.

Thus a second equilibrium, ballooning stable near the edge of the plasma, was calculated for the purposes of the present paper. This case has a more peaked pressure profile than the first, but both have average $\beta \approx 1\%$, where β is the ratio of plasma pressure to magnetic field pressure.

The q -profiles are not monotonic—in the peaked pressure profile case studied in this paper, q was 0.8895 on the magnetic axis, rising to a maximum value of 0.8964 quite close to the magnetic axis, then falling monotonically to 0.8675. Clearly the (global) magnetic shear is

very weak. Despite this fact and the non-monotonicity, there is some formal simplification in choosing $s \equiv q$, and we have taken $s = q$ since the region of plasma studied is in a monotonic-decreasing part of the q -profile (the decreasing region outside the maximum- q surface).

In these scans the most unstable ballooning eigenvalue was tabulated on a three-dimensional grid in s, α, θ_k space. The dependence on α was found to be rapid. The dependence on θ_k was much slower, but the variation was sufficient that the higher-growth-rate isosurfaces formed a set of distinct, topologically spherical branches. It was argued in [22] that this branch structure is produced by Anderson localization in bad curvature regions due to the strong breaking of both helical and axisymmetry in H-1NF.

According to the perturbation expansion in q' described in [22], a quadratic form in α, θ_k should form a good approximation to $\lambda - \lambda_{\min}(q)$ in the neighborhood of the central branch. Accordingly a least-squares fit on each surface was performed to provide a simple analytical description of the (0,0) [22] branch.

The radial dependence of the fitting coefficients was approximated by fitting to third-degree polynomials in q . Sections of the resulting approximation to the central branch are shown in Fig. 1. The isosurface spans a substantial range of magnetic surfaces within the plasma—the narrow range of variation in q is due to the low magnetic shear in H-1NF.

In order to establish the nature of the ray dynamics described by the regularized Hamiltonian, Eq. (4), a numerical integration with cutoff at $k_{\max} = 50$ was performed with initial conditions $q = q_2, \alpha = 0$, and $k_\alpha = 5$, where $[q_1, q_2] = [0.8852, 0.8951]$ is the q -range spanned by the $\lambda = -6$ isosurface as seen in Fig. 1. (A run with $k_\alpha = 10$ was also performed, with similar results.) Choosing the value $K = 1$ gave a good compromise between the sharp boundary potential to be modeled, and the smooth potential required for the numerical integration. The orbit remained on the “energy shell” $\lambda = -6$ to within an accuracy of one part in 10^6 over the “time” interval of the integration, 7500.




FIG. 2. Two views of intersections with the Poincaré surface of section $\alpha = 0$.

The two Poincaré plots in Fig. 2 show the orbit to be strongly chaotic, filling the “energy shell” ergodically, except that the regions $k_\alpha > 0$ and $k_\alpha < 0$ are dynamically

disjoint. The solid curve shown surrounding the outer limits of the “energetically accessible” region is calculated by solving $\lambda(0, q, k_q/k_{\max}) = -6$.

According to the Weyl formula [16, pp. 257–261], the number, $N(\lambda_{\max})$, of global eigenmodes with eigenvalues below the eigenvalue λ_{\max} is given, asymptotically in the limit $N \rightarrow \infty$, as $N(\lambda_{\max}) \sim v_{4D}(\lambda_{\max})/(2\pi)^2$. Here $v_{4D}(\lambda_{\max})$ is the volume of the dynamically accessible 4-dimensional phase-space region $\lambda(\alpha, q, k_q/k_{\alpha}) < \lambda_{\max}$, $0 < k_{\alpha} < k_{\max}$. The k_{α} integration can be performed analytically, giving $v_{4D}(\lambda_{\max}) = \frac{1}{2}k_{\max}^2 v_{3D}(\lambda_{\max})$, where $v_{3D}(\lambda_{\max})$ is the volume within the isosurface $\lambda(\alpha, q, \theta_k) = \lambda_{\max}$. Thus

$$N(\lambda_{\max}) \sim \frac{1}{8\pi^2} k_{\max}^2 v_{3D}(\lambda_{\max}). \quad (5)$$

We can make a rather rough estimate of the minimum value of n_{\max} required for CAS3D or TERPSICHOE to find even one eigenvalue with $\lambda < \lambda_{\max}$ by setting $N(\lambda_{\max}) = 1$ and calculating $k_{\max} \approx n_{\max}$ from Eq. (5). This gives $n_{\max}(N = 1) \sim (8\pi^2/v_{3D})^{1/2}$.

The isosurface $\lambda = -6$ studied above is about the largest of the disjoint topologically spherical isosurfaces corresponding to the highly toroidally localized strongly ballooning unstable regions of α, q, θ_k space. (For $\lambda > -6$ the isosurfaces are no longer topologically spherical.) Using the polynomial fits described above, we calculate $v_{3D}(-6) = 0.02158$. This gives $n_{\max}(N = 1) \approx 60$. Assuming that the dominant contributions to the MHD energy δW come from the rational surfaces intersecting the $\lambda = -6$ isosurface, we thus predict that it would be necessary to include, as a minimum set, basis functions having $(n, m) = (9, 8), (18, 16), (19, 17), (27, 24), (28, 25), (35, 31), (36, 32), (37, 33), (38, 34), (44, 39), (45, 40), (46, 41), (47, 42), (53, 47), (54, 48), (55, 49), (56, 50),$ and $(57, 51)$ to resolve a toroidally localized ballooning mode. (Here n, m are the toroidal and poloidal Fourier mode numbers, respectively.)

The large value of $n_{\max}(N = 1)$ required, and the unusual spread in n required in the basis set, will make these modes difficult to resolve using global eigenvalue codes (e.g. the simplifying phase factor method sometimes used in CAS3D studies [1] would not be appropriate). It is hoped that the Weyl formula estimate above will act as a guide in a future more extensive study using such a code. Physically, the large value of n_{\max} suggests that toroidally localized ballooning modes in H-1NF should be subject to strong FLR stabilization.

We can also apply the same approach to the toroidally localized ballooning branches found in the Large Helical Device (LHD) study [12]. From the plots in [12] we estimate $v_{3D} \sim 0.05$, which gives $n_{\max}(N = 1) \approx 40$.

The ballooning calculations were carried out on the Australian National University Supercomputer Facility’s Fujitsu VPP300 vector processor. We thank Dr. H. J. Gardner for providing the H-1 heliac VMEC input files

and Dr. S. P. Hirshman for use of the VMEC equilibrium code. Some of this work was done while one of us (RLD) was a visiting scientist at Princeton University Plasma Physics Laboratory, supported under US DOE contract No. DE-AC02-76CH0-3703. Useful conversations with Drs. M. Redi and A.H. Boozer are gratefully acknowledged.

-
- [1] A. H. Reiman *et al.*, Plasma Physics Reports **23**, 472 (1997).
 - [2] A. Reiman *et al.*, Plasma Phys. Control. Fusion **41**, B273 (1999).
 - [3] G. H. Nielson *et al.*, Phys. Plasmas **7**, 1911 (2000).
 - [4] B. D. Blackwell, Bull. Am. Phys. Soc. **45**, 289 (2000).
 - [5] I. B. Bernstein, E. A. Frieman, M. D. Kruskal, and R. M. Kulsrud, Proc. R. Soc. London Ser. A **244**, 17 (1958).
 - [6] R. L. Dewar, J. Plasma and Fusion Res. **73**, 1123 (1997).
 - [7] R. L. Dewar and A. H. Glasser, Phys. Fluids **26**, 3038 (1983).
 - [8] D. V. Anderson *et al.*, Int. J. Supercomp. Appl. **4**, 34 (1990).
 - [9] C. Schwab, Phys. Fluids B **5**, 3195 (1993).
 - [10] R. L. Dewar, J. Manickam, R. C. Grimm, and M. S. Chance, Nucl. Fusion **21**, 493 (1981), corrigendum: Nucl. Fusion, **22** (1982) 307.
 - [11] W. A. Cooper, D. B. Singleton, and R. L. Dewar, Phys. Plasmas **3**, 275 (1996), erratum: Phys. Plasmas **3**, 3520 (1996).
 - [12] P. Cuthbert *et al.*, Phys. Plasmas **5**, 2921 (1998).
 - [13] E. Hameiri, Commun. Pure Appl. Math. **38**, 43 (1985).
 - [14] A. E. Lifschitz, *Magnetohydrodynamics and Spectral Theory* (Kluwer, Dordrecht, The Netherlands, 1989), pp. 416–423.
 - [15] W. M. Nevins and L. D. Pearlstein, Phys. Fluids **31**, 1988 (1988).
 - [16] M. C. Gutzwiller, *Chaos in Classical and Quantum Mechanics, Interdisciplinary Applied Mathematics Series, Vol. 1* (Springer-Verlag, New York, 1990).
 - [17] E. Ott, *Chaos in Dynamical Systems* (Cambridge Univ. Press, Cambridge, U.K., 1993).
 - [18] W. M. Tang, R. L. Dewar, and J. Manickam, Nucl. Fusion **22**, 1079 (1982).
 - [19] R. J. Hastie, P. J. Catto, and J. J. Ramos, Bull. Am. Phys. Soc. **45**, 363 (2000).
 - [20] S. M. Hamberger, B. D. Blackwell, L. E. Sharp, and D. B. Shenton, Fusion Technol. **17**, 123 (1990).
 - [21] S. P. Hirshman and O. Betancourt, J. Comput. Phys. **96**, 99 (1991).
 - [22] P. Cuthbert and R. L. Dewar, Phys. Plasmas **7**, 2302 (2000).
 - [23] W. A. Cooper and H. J. Gardner, Nucl. Fusion **34**, 729 (1994).

The Princeton Plasma Physics Laboratory is operated
by Princeton University under contract
with the U.S. Department of Energy.

Information Services
Princeton Plasma Physics Laboratory
P.O. Box 451
Princeton, NJ 08543

Phone: 609-243-2750
Fax: 609-243-2751
e-mail: pppl_info@pppl.gov
Internet Address: <http://www.pppl.gov>