


PPPL-4189

PPPL-4189

**High Frequency Shear Alfvén Instability
driven by Circulating Energetic Ions
in NSTX**

Ya.I Kolesnichenko, R.B. White,
and Yu.V. Yakovenko

November 2006



Princeton Plasma Physics Laboratory

Report Disclaimers

Full Legal Disclaimer

This report was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor any agency thereof, nor any of their employees, nor any of their contractors, subcontractors or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or any third party's use or the results of such use of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

Trademark Disclaimer

Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof or its contractors or subcontractors.

PPPL Report Availability

Princeton Plasma Physics Laboratory:

http://www.pppl.gov/pub_report/

Office of Scientific and Technical Information (OSTI):

<http://www.osti.gov/bridge>

U.S. Department of Energy:

U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831-0062
Telephone: (865) 576-8401
Fax: (865) 576-5728
E-mail: reports@adonis.osti.gov

High frequency shear Alfvén instability driven by circulating energetic ions in NSTX

Ya. I. Kolesnichenko,¹ R. B. White,² and Yu. V. Yakovenko¹

¹*Institute for Nuclear Research, Prospect Nauky 47, Kyiv, 03680, Ukraine*

²*Princeton Plasma Physics Laboratory,*

P.O. Box 451, Princeton, New Jersey 08543, USA

(Dated: September 14, 2006)

Abstract

It is shown that a number of features of an instability with the frequency comparable to the ion gyrofrequency observed in the National Spherical Torus Experiment (*E. D. Fredrickson et al., Report P5.058 at the 33rd EPS Conference on Plasma Physics, Rome, 2006*) are consistent with the features of the Alfvén instability with large, about the inverse Larmor radius of the energetic ions (ρ_b^{-1}) longitudinal wave numbers. The conclusions drawn are based on an analysis of the resonant interaction of the energetic circulating ions and the waves, as well as on the calculation of the instability growth rate taking into account effects of the finite Larmor radius, ρ_b .

I. INTRODUCTION

In a pioneer work on Alfvén instabilities driven by the energetic ions an instability with the wave numbers along and across the magnetic field (k_{\parallel} and k_{\perp}) comparable to the inverse fast ion gyroradius, ρ_b^{-1} , was predicted.^{1,2} Later energetic-ion driven Alfvén instabilities were observed in many experiments on tokamaks, spherical tori, and stellarators, see, e.g., overviews in Refs. 3,4. However, in contrast to the instability considered in Ref. 1, all the reported instabilities observed experimentally had very small longitudinal wave numbers, $k_{\parallel} \ll \rho_b^{-1}$, and the frequencies which were much, by several orders, less than the ion gyrofrequency, ω_B . On the other hand, recently an instability of Alfvénic type with the higher frequencies was observed in the National Spherical Torus Experiment (NSTX).⁵ In the mentioned experiment, the frequencies at the beginning of the instability bursts were only by a factor of 5 less than the ion gyrofrequency. Therefore, it is natural to assume that the mechanism of destabilization of these waves differs from that of the conventional Alfvén instabilities by involving the cyclotron wave-particle interaction. This possibility is investigated below. A new Alfvén instability with the wave numbers about the energetic ion gyrofrequency is considered, which takes into account both effects of the finite Larmor radius of the energetic ions and the particle drift motion induced by the toroidicity. A comparison is made of the features of this instability and the features of the experimentally observed waves.

The structure of the work is as follows. In Sec. II we carry out a qualitative analysis of a possible resonant wave-particle interaction in an NSTX experiment. In Sec. III an expression for the instability growth rate is derived; it is used to evaluate the magnitude of the instability growth rate in the experiment and analyze the role of the anisotropy and the spatial distribution of the energetic ions.

II. QUALITATIVE ANALYSIS OF THE DESTABILIZATION OF HIGH-FREQUENCY ALFVÉN WAVES

We consider the case when the energetic ions are produced by the tangential neutral beam injection in the direction of the magnetic field, so that the energetic ion population

consists of circulating particles with

$$0 < v_{\parallel} < v_b, \quad (1)$$

where v_{\parallel} is the longitudinal velocity of the energetic ions and v_b is their birth velocity. These particles can lead to Alfvén instabilities through the resonant interaction with the waves. Typically, the destabilized waves have small longitudinal wave numbers, $k_{\parallel} \sim 1/qR$ (R is the major radius of the torus, q is the safety factor), and the wave frequencies, ω , negligible in the comparison with the ion cyclotron frequency, ω_B . However, we are interested in waves with ω comparable to ω_B . In this case $k_{\parallel}qR \gg 1$ and, therefore, the conventional sideband resonance, $\omega = [k_{\parallel} \pm 1/(qR)]v_{\parallel}$, responsible for instabilities with $k_{\parallel}qR \sim 1$ leads to the resonant velocity $v_{\parallel}^{res} \approx \sigma_k v_A$, with $\sigma_k = k_{\parallel}/|k_{\parallel}|$ and v_A the Alfvén velocity (we assume without loss of generality that $\omega > 0$). However, the efficiency of this resonance is rather low in systems with weak magnetic field because of the low ratio v_A/v_b (in the NSTX experiment considered in this work $v_A/v_b < 1/3$). Moreover, this resonance does not work for the waves propagating in the counter direction with respect to the beam, which was the case in the NSTX experiment. Therefore, we have to assume that the instability is associated with a resonance involving the gyrofrequency. The corresponding resonance condition can be written as

$$\omega = k_s \langle v_{\parallel} \rangle + l \langle \omega_B \rangle, \quad (2)$$

where $k_s = k_{\parallel} + s/(qR)$, s and l are integers, $\langle \dots \rangle = \oint dt(\dots)/\tau_b$, τ_b is the particle transit time. For Alfvén waves the longitudinal wave number is $k_{\parallel} \approx \sigma_k \omega/v_A$. Taking this into account and assuming $v_{\parallel}^{res} > v_A$, we obtain from Eqs. (1) and (2) that the considered injected ions and the waves can be in the resonance provided that the wave frequency is not less than a certain magnitude, ω_{\min} , taking place for $l = 1$ and given by

$$\omega_{\min} = \frac{\omega_B I_s}{v_b/v_A - \sigma_k}, \quad (3)$$

where $I_s = 1 - |s|\rho_b/(qR)$, $\rho_b = v_b/\omega_B$. Note that Eq. (3) was obtained in the assumption that it is mainly low- s resonances with $|s| < qR/\rho_b$ that contribute to the destabilization of the waves. For $v_b \gg v_A$ and $I_s \sim 1$, it follows from Eq. (3) that the longitudinal wave number of the waves interacting resonantly with the particles is $k_{\parallel} \sim \rho_b^{-1}$.

Equation (3) shows that the minimum frequency of the counter-propagating waves ($k_{\parallel} < 0$) is smaller than that of the co-propagating waves. In particular, for the NSTX experiment

with $v_b/v_A = 3.5$, $R = 100$ cm, $q = 2$, $v_b = 3 \times 10^8$ cm \cdot s $^{-1}$, $\omega_B = 1.4 \times 10^7$ s $^{-1}$ we obtain $\omega_{\min} \approx 450$ kHz for $k_{\parallel} < 0$ and $\omega_{\min} \approx 800$ kHz for $k_{\parallel} > 0$. These magnitudes take place when $s = -1$.

When the plasma rotates toroidally in the beam direction and $\text{sgn}(k_{\varphi}) = \text{sgn}(k_{\parallel})$ (φ is the toroidal angle), the Doppler effect decreases the frequency of counter-propagating waves and increases the frequency of co-propagating waves in the laboratory frame ($\omega^{\text{lab}} = \omega + \mathbf{k} \cdot \mathbf{u}$, with \mathbf{u} the velocity of the plasma rotation). As a result, the difference between ω_{\min} of co-propagating waves and ω_{\min} of counter-propagating waves increases. To evaluate this effect in the NSTX experiment, we take the toroidal mode number $n = 5$ and $u = 10^7$ cm \cdot s $^{-1}$. Then $k_{\varphi} = -n/R = -0.05$ cm $^{-1}$ [we take perturbations in the form $\propto \exp(-i\omega t + im\vartheta - in\varphi)$, where ϑ is the poloidal angle, m and n are the poloidal and toroidal mode numbers, respectively], and the Doppler shift is 80 kHz. This leads to $\omega_{\min}^{\text{lab}} = 370$ kHz for the counter-propagating waves and $\omega_{\min}^{\text{lab}} = 880$ kHz for the co-propagating waves.

This simple analysis enables us to conclude the following. First, the resonance (2) is consistent with the frequencies of the destabilized waves in the NSTX experiment (in particular, the frequencies at the beginning of the instability bursts were about 400 kHz in the shot #114147 and 500 kHz in the shot #114154). Second, only counter-propagating waves can be destabilized in the frequency range relevant to the experiment. Third, the resonance condition imposes restrictions on the instability mode numbers. They are $n > 0$ and $m < nq$, which follows from two conditions, $k_{\varphi} = -n/R < 0$ found experimentally, and $k_{\parallel} = (m - nq)/(qR) < 0$ required for the frequency of $\sim 400 - 500$ kHz to satisfy the resonance condition. The obtained restriction on the poloidal mode numbers is valuable because it may help to select the mode among those which can be found by numerical simulation (the poloidal mode numbers are not available from experimental measurements). The restriction on n is also important in connection with the following.

The fulfillment of the resonance condition is not sufficient for the instability to arise. The factors which can lead to the destabilization of the waves are spatial inhomogeneity of the energetic ions and their velocity anisotropy. The spatial inhomogeneity drives the instability when $n dn_b/dr > 0$ (n_b is the beam density). Because $n > 0$, the beam particles with monotonically decreasing radial profile have a stabilizing (rather than destabilizing) influence on the waves in the case of counter-rotating modes. However, calculations show, that in the NSTX case the beam radial distribution has an off-axis maximum, which suggests

that the beam spatial inhomogeneity tends to destabilize modes localized in the core region. A more detailed analysis is required to clarify the role of the velocity anisotropy. This will be done in the next section.

III. CALCULATION OF THE INSTABILITY GROWTH RATE

In order to evaluate the instability growth rate (γ) we use the local approximation. Neglecting the wave damping caused by the bulk plasma we can write

$$\frac{\gamma}{\omega} = -\frac{v_A^2}{2c^2}\epsilon_1, \quad (4)$$

where ϵ_1 is the fast ion dielectric permeability tensor component calculated in Ref. 6. We take into account that $F = F(r, \mathcal{E}, \lambda, \sigma_v)$, where F is the unperturbed distribution function of the energetic ions, $\int d^3v F = n_b$, \mathcal{E} is the particle energy, $\lambda = \mu B_0/\mathcal{E}$, μ is the particle magnetic moment, B_0 is the magnetic field at the magnetic axis, $\sigma_v = \text{sgn}(v_{\parallel})$. Using the resonance condition given by Eq. (2), we calculate the integral over the energy in the expression for ϵ_1 . Then we obtain:

$$\frac{\gamma}{\omega} = \frac{\pi^2 \omega_B^4}{k_{\perp}^2 n_i \omega^2} \sum_{l,s,\sigma_v} \frac{l^2}{|k_s|} \int \frac{d\lambda}{1-\lambda} J_l^2(\xi) J_s^2(\zeta) \hat{\Pi} F \Big|_{v^{\text{res}}}, \quad (5)$$

where n_i is the bulk ion density, $v^{\text{res}} = (\omega - l\omega_B)/(k_s \sqrt{1-\lambda})$ is the resonant velocity, $J_l(\xi)$ and $J_s(\zeta)$ are the Bessel functions, $\xi = k_{\perp} \rho \sqrt{\lambda}$, $\zeta = (q/\sqrt{1-\lambda}) \sqrt{k_{\perp}^2 \rho^2 (1-0.5\lambda)^2 / \kappa^2 + l^2 r^2 / \rho^2}$, $\rho = v/\omega_B$, k_{\perp} is the transverse wave number, κ is the plasma elongation,

$$\hat{\Pi} = \omega \frac{\partial}{\partial \mathcal{E}} + (l\omega_B - \lambda\omega) \frac{1}{\mathcal{E}} \frac{\partial}{\partial \lambda} + \frac{nq}{\omega_B M} \frac{1}{r} \frac{\partial}{\partial r}. \quad (6)$$

Note that Eq. (5) takes into account effects of finite transverse Larmor radius of the energetic ions ($\rho_{\perp} = \rho \sqrt{\lambda}$). Moreover, the growth rate is considerable only when $k_{\perp} \rho_{\perp}$ is not small.

Equation (5) can be used to calculate the growth rate numerically. We, however, restrict ourselves to an analytical evaluation of γ . With this purpose we approximate the distribution function of the energetic ions as follows:

$$F = n_b f_{\mathcal{E}} f_{\lambda} \eta(v_b - v) \eta(\sigma_v), \quad (7)$$

with $f_\xi = C/\mathcal{E}^{3/2}$, $C^{-1} = M^{-3/2}\pi\sqrt{2}\Lambda \int d\lambda f_\lambda/\sqrt{1-\lambda}$ is a constant found from the equation $\int d^3v f_\xi = 1$, $\Lambda = \ln v_b/v_c$ with $v_c \sim (M_i/M_e)^{1/3}$, $f_\lambda = f_\lambda(\lambda)$, $\int f_\lambda d\lambda = 1$, $\eta(x) = \int_{-\infty}^x \delta(x)dx$. For the considered waves the first term in Eq. (6) can be neglected. The term associated with the velocity anisotropy [proportional to the λ -derivative in Eq. (6)] can be transformed by integrating by parts. Assuming that particles with small λ dominate in the energetic-ion population, we neglect for simplicity the terms proportional to $\partial J_s(\zeta)/\partial\lambda$ and use the approximation $d\xi(v^{\text{res}})/d\lambda = \xi^{\text{res}}/(2\lambda)$. Then we obtain:

$$\frac{\gamma}{\omega} = \frac{\pi n_b \omega_B^2}{\Lambda n_i \omega^2} \left(\int \frac{d\lambda f_\lambda}{\sqrt{1-\lambda}} \right)^{-1} \times \sum_{l,s} \frac{l^2}{|l - \omega/\omega_B|} \int \frac{d\lambda f_\lambda}{\sqrt{1-\lambda}} J_s^2(\zeta) \left[-\frac{l}{\xi} \frac{\partial J_l^2(\xi)}{\partial \xi} + J_l^2(\xi) \frac{nq}{k_\perp^2 r} \frac{\partial \ln n_b}{\partial r} \right]_{v^{\text{res}}}. \quad (8)$$

In the case relevant to the NSTX experiment, only the $l = +1$ harmonic contribute (there are no resonant particles for other l). Therefore, the velocity anisotropy of the energetic ion distribution is a destabilizing factor only when the particle Larmor radius is sufficiently large to lead to negative derivative of $J_l^2(\xi)$. One can see that this condition is satisfied even for a low- m instability when the mode is well localized. For $f_\lambda \propto \delta(\lambda - \lambda_0)$ the term associated with the anisotropy dominates, unless ξ is about a point where $dJ_l^2/d\xi = 0$.

Let us evaluate the growth rate in the NSTX shot #114147. We take $n_b/n_i \sim 1/20$, $\omega_B/\omega = 4.7$, $\zeta \sim 10$, $\xi \sim 2$ and $J_n^2(x) \sim (\pi x)^{-1}$. Then we obtain $\gamma/\omega \sim (\pi\zeta\xi^2)^{-1}$, which leads to $\gamma/\omega \sim 10^{-2}$ for $\xi \sim 2$ and $\zeta \sim 10$. Numerical calculations with using Eq. (5) give a smaller growth rate because in reality the beam distribution function is not peaked at very small λ .

Thus, the driving part of the instability growth rate is rather small. Therefore, the perturbative approach is justified in spite of the fact that the beam pressure was comparable to the plasma pressure in the core region in the considered NSTX shot.

IV. SUMMARY AND CONCLUSIONS

Our analysis shows that the destabilized waves with the frequencies $\omega \sim \omega_B/5$ observed in NSTX in the shots #114147 and #114154 during neutral beam injection can be identified as the high frequency Alfvén instability with the $k_\parallel \rho_b \sim 1$. This instability arises due to the cyclotron resonant interaction of Alfvén waves and the beam particles. The following facts

support this conclusion. First, the wave frequencies determined by the resonance condition are close to the observed frequencies. Second, our analysis predicts that only counter-propagating (with respect to the beam direction) waves can be destabilized, in agreement with the experiment. Third, the resonance condition leads to the toroidal mode number $n > 0$, again in agreement with the experiment. In addition, it imposes a restriction on the poloidal mode number, $m < nq$. The considered instability arises mainly because of the velocity anisotropy of the injected ions, although the spatial inhomogeneity – the hollow radial distribution of the energetic ions – contributes to the destabilization of the waves with $n > 0$. The instability growth rate (the driving part) is relatively small, which justifies the use of the perturbative approach in the analysis.

Acknowledgments

This work was supported in part by the Award No. UKP2-2643-KV-05 of the U.S. Civilian Research & Development Foundation, and by the U.S. Department of Energy Grant DE-FG03-94ER54271. The authors thank E. D. Fredrickson and N. Gorelenkov for providing experimental data and discussions. Two authors (Y.K. and Y.Y.) would like to acknowledge the hospitality of the Princeton Plasma Physics Laboratory.

-
- ¹ V. S. Belikov, Ya. I. Kolesnichenko, and V. N. Oraevskij, *Zh. Ehksp. Teor. Fiz.* **55**, 2210 (1968); *Sov. Phys. – JETP* **28**, 1172 (1969).
- ² Ya. I. Kolesnichenko, *Nucl. Fusion* **20**, 727 (1980).
- ³ K. L. Wong, *Plasma Phys. Control. Fusion* **41**, R1 (1999).
- ⁴ A. Weller, M. Anton, J. Geiger *et al.*, *Phys. Plasmas* **8**, 931 (2001).
- ⁵ E. D. Fredrickson, N. Gorelenkov, and J. Menard, “Observation of Hole-Clump Pair Generation by Global or Compressional Alfvén Eigenmodes”, Proceedings of the 33rd European Physical Society Conference on Plasma Physics, Rome, 2006, Report P5.058, Europhysics Conference Abstracts, to be published.
- ⁶ V. S. Belikov, Ya. I. Kolesnichenko, and R. B. White, *Phys. Plasmas* **10**, 4771 (2003).

The Princeton Plasma Physics Laboratory is operated
by Princeton University under contract
with the U.S. Department of Energy.

Information Services
Princeton Plasma Physics Laboratory
P.O. Box 451
Princeton, NJ 08543

Phone: 609-243-2750
Fax: 609-243-2751
e-mail: pppl_info@pppl.gov
Internet Address: <http://www.pppl.gov>